

Identification of genes expressed during the compatible interaction of grapevine with *Plasmopara viticola* through suppression subtractive hybridization (SSH)

Guillaume Legay · Elaheh Marouf · Dave Berger · Jean-Marc Neuhaus · Brigitte Mauch-Mani · Ana Slaughter

Abstract Grapevine (*Vitis vinifera*) is the most widely cultivated and economically important fruit crop, but is susceptible to a large number of diseases. Downy mildew, caused by the obligate biotrophic oomycete pathogen *Plasmopara viticola*, is a common disease present in all regions where vines are cultivated. We used suppression subtractive hybridization (SSH) to generate two cDNA libraries enriched for transcripts induced and repressed, respectively, in the susceptible grapevine cultivar Chasselas 24 h after inoculation with *P. viticola*. Differential screening on glass slide microarrays yielded over 800 putative genes that were up-regulated in response to *P. viticola* infection and over 200 that were down-regulated. One hundred and ninety four of these, were sequenced, identified and functionally categorised. Transcript abundance of twelve genes over a 48 h time course was examined

by reverse transcriptase quantitative real-time PCR (RT-qPCR). Ten of these genes were induced/enhanced by *P. viticola* challenge, confirming the results of the SSH. The vast majority of the genes identified are related to defence. Interestingly, many genes involved in photosynthesis were down-regulated.

Keywords Compatible interaction · Reverse transcriptase quantitative real-time PCR · *Plasmopara viticola* · Suppression subtractive hybridization · *Vitis vinifera*

Introduction

Downy mildew caused by the obligate biotrophic pathogen *Plasmopara viticola*, is an economically important disease of grapevine. The pathogen attacks all green parts of the grapevine. Within a few hours after inoculation, the pathogen has penetrated the tissue and formed the first haustoria in a susceptible cultivar and after 3 days, the intercellular spaces are entirely filled with mycelium and sporulation is abundant under favourable environmental conditions (Unger et al. 2007). In a resistant cultivar, the first infection steps are the same, however, the invasive growth of *P. viticola* is delayed and further development ceases before the intercellular spaces colonized (Unger et al. 2007). Most of the widely grown grapevine cultivars are highly susceptible to *P.*

G. Legay · E. Marouf · J.-M. Neuhaus ·
B. Mauch-Mani (✉) · A. Slaughter
Laboratory of Molecular and Cellular Biology,
University of Neuchâtel,
Rue Emile-Argand 11, Case Postal 158, 2009 Neuchâtel,
Switzerland
e-mail: Brigitte.mauch@unine.ch

D. Berger
Department of Plant Science, Forestry and Agricultural
Biotechnology Institute (FABI), University of Pretoria,
Lunnun Road,
Pretoria 0002, South Africa

viticola and the control of downy mildew requires regular fungicide applications. Application of copper-containing fungicides to control downy mildew causes accumulation of this heavy metal in soil and groundwater, resulting in toxic effects to the environment. Constant use of fungicides is also a problem since it may favour the emergence of fungicide-resistant isolate that may dominate the population.

Although a number of studies describe the biology of the infection process of grapevine by *P. viticola* (Allegre et al. 2007; Kortekamp 2005; Musetti et al. 2007; Unger et al. 2007; Werner et al. 2002) and the characterisation of defence reactions in susceptible and resistant grapevines (Dai et al. 1995; Gindro et al. 2003; Kortekamp 2006, Kortekamp and Zyprian 2003) or their response to inducers of defence reactions (Aziz et al. 2003; Hamiduzzaman et al. 2005; Trouvelot et al. 2008), a large-scale analysis of *P. viticola* and grapevine interaction has not been reported.

Upon recognition of an invader, signal transduction events ultimately leading to the induction of both chemical and physical defence responses are activated (Scheel 1998). The specific recognition of a pathogen often leads to a programmed cell death termed the hypersensitive response (HR). It is a rapid and efficient plant resistance mechanism leading to cell death at the site of infection, usually in gene-for-gene interactions with obligate biotrophic pathogens (Heath 2000). The cellular events taking place during the HR comprise among others the generation of reactive oxygen species (ROS) at the site of infection, cell wall reinforcements through callose-rich deposits, ion fluxes, and the synthesis of anti-microbial compounds and enzyme inhibitors (Glazebrook 2005). In grapevine, these include low-molecular antimicrobial compounds (phytoalexins), deposition of phenolics, lignin and callose and increased activity of pathogenesis-related (PR) proteins with hydrolytic activity (chitinases and glucanases; Derckel et al. 1999).

Since the difference between a resistant and susceptible plant is often just a question of timing and amplitude of the adequate defence response (Polesani et al. 2010), investigating a compatible interaction not only gives information on the availability of adequate defence mechanisms but will also assist in the development of new control strategies and lead to the identification of pathogen and host factors needed for disease progression. Investigating the molecular basis of plant-pathogen interactions often uncovers novel aspects of plant cell

biology and signalling mechanisms. Expression profiles investigating compatible interactions in cultivated grapevine genotypes is limited and downy mildew has received very little attention compared to diseases caused by other biotrophic pathogens, such as powdery mildew and rusts (Polesani et al. 2008).

Functional genomic approaches provide powerful tools for identifying expressed genes. Among these techniques, expressed sequence tags (EST) (Adams et al. 1991), serial analysis of gene expression (SAGE) (Velculescu et al. 1995) and massively parallel signature sequencing (MPSS) (Brenner et al. 2000), have been successfully employed. Subtractive suppression hybridization (SSH) is an effective method that can be used to maximise the identification of genes that are involved in host responses to pathogen infection and disease development. This technique has been used to isolate plant genes that are expressed in response to infection (Birch et al. 1999; Degenhardt et al. 2005; Lu et al. 2004; Van den Berg et al. 2007). The SSH cDNA library approach reduces the cloning of abundantly expressed housekeeping genes or genes commonly expressed in both control and treated plants, therefore significantly enhancing the chances of cloning differentially expressed genes. This is particularly important because many PR genes are expressed at low levels. These genes are less likely to be represented in a library if standard EST cloning methods are used. The aim of this study was to construct a SSH cDNA-library to identify and verify genes that are differentially expressed during a susceptible response of *V. vinifera* cv. Chasselas to *P. viticola*. By this approach, *P. viticola*-responsive genes were identified and classified into functional groups and the expression profiles of some of the genes were examined by quantitative real time PCR following inoculation with *P. viticola*.

Material and methods

Biological materials

Grapevine plants were grown from *V. vinifera* cv Chasselas seeds (susceptible to *P. viticola*), in soil at Syngenta (Stein, Switzerland) in a glasshouse at day/night temperature of 23°C/ 18°C with a photoperiod of 16 h and relative humidity (RH) of 65±10%. Seedlings at the five-to six-leaf stage (6 weeks) were used for infection experiments.

Pathogen and inoculation of plants

P. viticola (house isolate, obtained from Syngenta, Stein, Switzerland) was maintained on susceptible Chasselas seedlings in a glasshouse. Weekly, the abaxial surfaces of young leaves were inoculated with an aqueous suspension containing $\sim 4 \times 10^4$ sporangia ml^{-1} . The inoculated plants were kept at 100% RH overnight. After incubation for 6 days under ambient glasshouse conditions, the plants were again maintained overnight at 100% RH to induce sporulation. The sporangia were harvested and used for inoculation.

Six week old grapevine seedlings were inoculated at Syngenta by spraying a freshly prepared sporangia suspension (6×10^4 sporangia ml^{-1}) onto the abaxial leaf surfaces. For the mock-inoculation, seedlings were sprayed with water. Inoculated and mock-inoculated seedlings were placed overnight in a humid chamber. Samples were collected from inoculated and mock-inoculated leaves 24 h after inoculation (for library construction) or at 0, 8, 12, 24 and 48 h after inoculation (for quantitative RT-PCR) and immediately frozen in liquid nitrogen and stored at -80°C . For the next 6 days the plants were placed in a glasshouse under the conditions described above and finally placed in a humid chamber overnight to induce sporulation. On day 7 plants were assessed for visual disease symptoms (Fig. 4a–c). For the library construction, 15 inoculated and 15 mock-inoculated plants were used. Twenty 4 h after inoculation, the third, fourth and fifth leaves from inoculated and mock-inoculated plants were harvested and immediately frozen in liquid nitrogen and stored at -80°C .

For gene expression analysis, at each time point a pool of 3 inoculated and 3 mock-inoculated plants was used. As for the library construction, the third, fourth and fifth leaves from the bottom of inoculated and mock-inoculated plants were harvested. Two independent experiments were carried out.

RNA isolation and SSH library construction

Total RNA was isolated from frozen leaf tissues using a modified CTAB extraction and lithium chloride precipitation method according to (Iandolino et al. 2004). The mRNA was isolated using Poly(A)Purist™ mRNA Purification Kit (Ambion) according to the manufacturer's protocol and purified mRNA quality was then determined on an Agilent 2100

Bioanalyser RNA 6000 Pico LabChip. Two micrograms of high quality mRNA were used for the construction of the libraries.

SSH was carried out using the PCR-Select cDNA Subtraction Kit (Clontech, Palo Alto, CA). cDNA prepared from the inoculated samples was used as the tester and that from the mock-inoculated samples as the driver for the forward subtraction to isolate fragments corresponding to genes whose expression level was increased following infection. The reverse library was carried out with the mock-inoculated samples as tester and inoculated samples as driver in order to isolate fragments corresponding to genes whose expression level decreased following infection. An Advantage PCR cloning kit (Clontech, USA) was used to selectively amplify the cDNA fragments preferentially present in the tester from the subtraction hybridization products. Secondary PCR products from the forward and reverse subtracted libraries were cloned into pGEM-T Easy vector (Promega) according to the manufacturer's protocol. Transformation was performed using Library Efficiency DH5 α Competent Cells (Invitrogen). A total of 1344 clones from the forward SSH library and 1152 from the reverse SSH library were randomly picked and stored in 96-well plates.

SSH cDNA microarray screening

In preparation for screening with microarrays, following the method of Van den Berg et al. (2004) and modifications of Berger et al. (2007), PCR reaction products from all the clones in the forward and reverse SSH libraries were purified using the 96-well Multiscreen HTS, FB plates (Millipore). An aliquot of the purified PCR products were visualised on a 1% agarose Electro-Fast Stretch gel to check that a single band was amplified and to estimate the yield. The purified PCR products were dried in a vacuum centrifuge and resuspended in 20 μl 3X SSC, 1.5 M betaine. Libraries containing in total 2496 grapevine clones were arrayed in triplicate onto aldehydesilane-coated slides (Nexterion™ Slide AL, Schott Nexterion, Jena, Germany) using an Omnigrid 300 contact-printing robotic microarrayer (Genomic Solutions, Ann Arbor, MI) equipped with SMP3 pins (TeleChem International Inc.). Spike controls (Lucidea Universal Scorecard, GE Healthcare) were included for each sub-grid of the microarray. Spot and printing quality were

assessed visually after printing and the DNA was cross-linked to the slides by baking at 80°C for 1 h. The slides were post-processed with sodium borohydride using the protocol recommended by the manufacturer (Schott Nexterion, Jena, Germany).

Preparation of fluorescent targets and hybridization

Forward subtracted (FS), forward unsorted (FU) and reverse unsorted (RU) cDNA targets were used to screen the forward SSH library, whereas reverse subtracted (RS), reverse unsorted (RU) and forward unsorted (FU) cDNA targets were used to screen the reverse SSH library on glass slide microarrays (Berger et al. 2007). cDNA targets of FS, FU, RS and RU, with the adaptors removed were labelled by incorporation of Cy5 and Cy3-dUTP using Klenow enzyme according to Berger et al. (2007). The labelled targets were purified using the Wizard SV gel and PCR clean-up system (Promega) and quantified using a Nanodrop spectrophotometer.

Eight slides were arrayed with the cDNA probes from the forward and reverse library separately. Before hybridization, equimolar amounts (300 pmol) of each pair of appropriate labelled targets and 10 µg yeast tRNA were mixed and concentrated through a Microcon YM-30 filter (Millipore). The target combinations for screening of the forward library were as follows: Cy3-labelled FS and Cy5-labelled FU; dye swap: Cy3-labelled FU and Cy5-labelled FS (to calculate Enrichment ratio 2); Cy3-labelled FU and Cy5-labelled RU; dye swap: Cy3-labelled RU and Cy5-labelled FU (to calculate Enrichment ratio 3) (Berger et al. 2007). For the screening of the reverse library, the target combinations were: Cy3-labelled RS and Cy5-labelled RU; dye swap: Cy3-labelled RU and Cy5-labelled RS (to calculate Enrichment ratio 2); Cy3-labelled RU and Cy5-labelled FU; dye swap: Cy3-labelled FU and Cy5-labelled RU (to calculate Enrichment ratio 3). For hybridizations, the labelled probes were combined with 3X SSC and 0.4% SDS in a final volume of 30 µl. The solution was boiled for 1 min, centrifuged and applied to the microarray under a cover slip. The microarray slides were sealed in a hybridization chamber (TeleChem) and submerged in a water bath at 64°C overnight. After hybridization, slides were washed twice for 5 min in 2X SSC/ 0.1% SDS, twice for 1 min in 0.2X SSC and twice for 1 min in 0.1X SSC. Slides were dried by centrifugation at 900 g for 2 min and scanned with an Agilent DNA

microarray scanner (Agilent Technologies). Genepix Pro 5.1 was used to localise and integrate every spot on the array. Analysis of microarray data was done according to Berger et al. (2007) by using the SSHscreen 1.0.4 package downloaded from <http://microarray.up.ac.za/SSHscreen/>. SSHscreen was used to implement the following statistical approaches: the normexp method was used to correct for the background, in order to normalise within arrays the printploess method was used and finally to normalise between arrays the Aquantile method was used.

Sequence analysis

Sequencing of the inserted cDNA fragments was done using the T7 and SP6 primers at Microsynth (Switzerland). Sequences were compared to the UniProt database as well as NCBI database using BLAST. Functional categorisation of sequences was performed by comparison with sequences in the NCBI database as well as through the use of the MIPS functional catalogue database (FunCatDB). Grapevine cDNA sequences have been deposited in GenBank dbEST with the accession numbers GO652854-GO653046.

Reverse transcriptase quantitative real-time PCR (RT-qPCR) analysis

To remove any contaminating DNA in the RNA preparations, RNA was treated with TURBO DNA-free DNase I (Ambion) according to the manufacturer's protocol. After DNase treatment, the RNA concentration and quality was checked with the NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). cDNAs were synthesised from 1 µg of total RNA using the Superscript III first strand synthesis system (Invitrogen) according to the manufacturer's instructions.

Real-time PCR reactions were performed in 96-well plates with a BioRad iQ iCycler Detection system (BioRad Laboratories, Ltd) using SYBR Green to detect dsDNA synthesis. The amount of transcripts of selected genes in *P. viticola*-inoculated versus mock-inoculated grapevine plants was compared by using the reference gene Elongation Factor 1- α (*EF1- α*) as endogenous reference for data normalization. Reactions were done in a total volume of 20 µl containing 200 nM of each primer, 2 µl of

1:5 diluted cDNA and 10 μ l 2X SYBR Green Master Mix Reagent (Bio-Rad). Reactions were run using the following cycling parameters: 95°C for 3 min, 40 cycles of 95°C for 30 s, 57°C for 35 s and 72°C for 35 s followed by a melt cycle from 60°C to 95°C. No-template controls were included for each primer pair to assess its specificity and each PCR reaction was done in triplicate. Primer sequences (Table 1) were designed with Primer Design 4 based on the sequences of the selected clones. The software programme Gene-X was used to calculate the mean normalised expression of the genes (Vandesompele et al. 2002).

Microscopic observation

Leaf disks from *P. viticola*-inoculated and mock-inoculated grapevine plants were collected at two time points (4 and 7 days). Leaf disks were then used for different staining methods.

Lactophenol-trypan blue staining

To observe the structure of *P. viticola*, leaf disks were stained with lactophenol trypan blue according to (Keogh et al. 1980). The disks were incubated for 48 h in the staining solution and then destained with chloral hydrate (2.5 gml⁻¹). Destained disks were kept in chloral hydrate solution and visualized with a light microscope. Hyphae, sporangiophores and sporangia stained blue.

Lignin staining

Leaf disks were boiled for 10 min in 95% ethanol and then treated with 10% phloroglucinol for 20 min and washed in 25% HCl. They were then mounted in glycerol (75%) and examined with a light microscope (Dai et al. 1995).

Flavonoid staining

Staining for the presence of flavonoids was performed according to (Pina and Errea 2008). Briefly, decolorized leaf disks were stained for 10 min in 1% (w/v) Naturstoff reagent A (2-Aminoethyl diphenylborinate, Fluka) in ethanol and then washed 4 times with ethanol. Visualization was done under an epifluorescence microscope with an UV filter (BP, 340–380 nm; LP, 450 nm)

Results

Construction of the SSH library

For the construction of cDNA libraries enriched for *P. viticola* responsive genes, pooled Chasselas plants inoculated with *P. viticola* or mock-treated plants were used for the extraction of mRNA and cDNA synthesis. Reciprocal subtractions of cDNAs derived from *P. viticola*-infected and non-infected plants yielded two cDNA libraries, one being enriched for *P. viticola*-responsive genes (forward library) and the other enriched for *P. viticola*-repressed genes (reverse library). In total 1344 clones form the forward SSH library and 1152 from the reverse SSH library were used.

Screening of the SSH library using glass slide microarrays

According to the PCR-select differential screening kit user manual (Clontech), 1–4% of the clones identified by differential screening with inverse dot blots, turn out to be false positives. In order to select clones for sequencing and reduce the number of false positives, a more quantitative, efficient, rapid and high-throughput technique was used, namely screening the forward and reverse SSH library on glass slide microarrays (as described in Van den Berg et al. 2004). In addition an analysis R software package named SSHscreen was used, which employs functions of limma (linear models for microarray data) to analyse spot intensity data, thereby screening clones in the libraries to identify those that are significantly differentially expressed (Berger et al. 2007; Coetzer et al. 2010). Furthermore, this method is able to identify whether the clones were derived from rare or abundant transcripts in the treated sample. Quantitative screening was carried out by calculating different “SSH enrichment ratios (ER)” for each clone. Calculations were based on the intensity values of each cDNA clone as a result of hybridization by the Cy dye-labelled cDNA targets. After hybridization, scanning and normalization of the data to account for intensity biases within and between slides, the SSH enrichment ratios were calculated and a statistical test was applied with a false discovery rate (FDR) of 5% to account for multiple testing. SSH enrichment ratio 3 (ER3) values were calculated as $\log_2(\text{FU}/\text{RU})$ (for

Table 1 Primers used for quantitative real-time RT-PCR

Gene name	Forward primer	Reverse primer
<i>Chalcone isomerase (CHI)</i>	5'-AACTTCTGGTAGGGACCCATCT-3'	5'-GAAGGATGAAACCTTCCCACCA-3'
<i>Chalcone synthase (CHS)</i>	5'-GAGAGCAACTCGACACGTTCTG-3'	5'-GTGTAGCAAGGCTGTGCAACAA-3'
<i>Chitinase III (Chit III)</i>	5'-ATCGTCTCGGCCATTAGGTGAT-3'	5'-ATACCTTCCTTCCACGCTTGCT-3'
<i>Cinnamate-4-hydroxylase (C4H)</i>	5'-AGTCCAAGTCACCGAGCCTGAT-3'	5'-TAGCAAAGCCACCATGCGTTCAC-3'
<i>EF1-α</i>	5'-GAACGTTGCTGTGAAGGATCTC-3'	5'-CGCCTGTCAACCTTGGTCAGTA-3'
<i>Flavonone-3-hydroxylase (F3H)</i>	5'-GAGAGCAACTCGACACGTTCTG-3'	5'-CTGAAGGAGCAGCGTGTGTT-3'
<i>β-1,3-Glucanase (Gluc)</i>	5'-CCAGTACAGCATCCGAAGAATGAG-3'	5'-AACCAGCAACGTATCGGAATCTGAC-3'
<i>Heavy-metal associated protein (HMAP)</i>	5'-CATCCACTGCCATGGAGCTT-3'	5'-CTTCTGCCACCGACCTAATC-3'
<i>Metallothionein (MT)</i>	5'-TGCATGGATCGCAGTTGCAGTT-3'	5'-TCGCACCAGTGAAGATGCACT-3'
<i>PR-1</i>	5'-AACGACACTGTAGCTGCCTACG-3'	5'-GCATCGGTGCCTGTCAATGAAC-3'
<i>PR-5</i>	5'-TGTTGTCGTGGGATCTCACCTG-3'	5'-CTTCAAAGGGTCAGGCAAGTGT-3'
<i>PR-10</i>	5'-CCAACCAATCCTCTTGTCTT-3'	5'-GAATCGAGGATAGAGGCCTAGC-3'
<i>PR-17</i>	5'-ATCGCCTGAGTAGCTTCCGATG-3'	5'-CGCTAGTATCTGCCACCGACTT-3'

the forward library) and $\log_2(\text{RU}/\text{FU})$ (for the reverse library) and gave an indication whether cDNAs were up- or down-regulated by *Plasmopara* infection (Fig. 1).

From the forward library screening, 441 grapevine clones were significantly up-regulated in response to *Plasmopara* infection with a positive ER3 value (and positive B statistics at FDR of 5%) and are represented in the top two quadrants of Fig. 1a. A number of clones in the forward library (161) had negative ER3 values and thus fell in the bottom two quadrants of Fig. 1a, indicating that they are “down-regulated” by *Plasmopara* infection, which means they are false positives and thus escaped the subtraction. SSHscreen produces a similar plot from the screening of the reverse library, but in this case positive ER3 values indicate grapevine clones that were down-regulated in response to *Plasmopara* infection (Fig. 1b). Screening of the reverse library resulted in a total of 64 clones that were down-regulated by *Plasmopara* infection (positive ER3 values, see top two quadrants of Fig. 1b) whereas 418 clones had negative ER values (i.e. escaped subtraction in the reverse library Fig. 1b). These results indicate that the efficiency of the SSH process was far better in the forward library compared to that of the reverse library (73% vs. 13%). The SSHscreen analysis thus enabled the selection of clones for DNA sequencing out of the 441 forward library clones that were scored as up-regulated, and

out of the 64 reverse library clones that were scored as down-regulated.

One of the main advantages of SSH is that it normalizes the cDNA abundance so that cDNAs encoded by genes that are expressed at low levels, but are nonetheless differentially regulated, can be identified readily. SSH enrichment ratio 2 (ER2) values were calculated as $\log_2(\text{FS}/\text{FU})$ (for the forward library) and $\log_2(\text{RS}/\text{RU})$ (for the reverse library) to give an indication whether cDNAs have been enriched or reduced relative to levels in the tester sample by the normalization process. Rare transcripts in *Plasmopara* infected plants will have negative inverse ER2 values and fall in the top left quadrant of Fig. 1a, while abundant transcripts will fall in the top right quadrant (Fig. 1a). In the forward library, half of the grapevine clones up-regulated by *Plasmopara* infection were rare transcripts, and half were abundant (48% vs. 52%). In the reverse library, the majority (75%) of the grapevine clones down-regulated by *Plasmopara* infection were rare transcripts (top left quadrant of Fig. 1b), while only 25% were abundant.

Sequencing analysis and functional classification of differentially expressed clones

No *P. viticola* genes were identified since the time point chosen to construct the SSH library was early,




Fig. 1 Screening of forward (a) and reverse (b) SSH grapevine libraries on glass slide microarrays. An up/down regulation plot of enrichment ratio 3 (ER3) vs. inverse enrichment ratio 2 (ER2) is produced by the SSHscreen analysis. The clones that were calculated in SSHscreen to be statistically significantly up-regulated (ER3 > 0, positive B statistic, false discovery rate of 5%) after *P. viticola* inoculation in the forward library (a) are shown in the top two quadrants, while those that are down-regulated in the forward library (false positives) are shown in the bottom two quadrants. The inverse is true for the reverse library (b), where the top two quadrants contain clones that are significantly down-regulated after *P. viticola* inoculation, and the bottom two quadrants contain false positive clones that are up-regulated. The inverse ER2 values reflect the relative abundance of the transcripts for each clone in the unsubtracted cDNA samples. Thus transcripts that were rare in the cDNA

from plants inoculated with *P. viticola*, but also were up-regulated relative to control plants fall in the top left quadrant (a), whereas those that were abundant in the cDNA from plants inoculated with *P. viticola*, but also were up-regulated relative to control plants fall in the top right quadrant (a). For the reverse library, the top left and top right quadrants show transcripts that were down-regulated after inoculation with *P. viticola* and that were rare and abundant, respectively, in the control plants prior to subtraction (b). No clones are observed in the region of the x-axis (ER3 = 0), since this region would contain genes that have similar transcript levels in inoculated and mock-inoculated samples, that would have been removed by the subtraction or not plotted on Fig. 1a and b since they are not calculated to be differentially expressed when applying a FDR of 5%

so that the amount of pathogen biomass (RNA) was very low compared to plant biomass.

A total of 223 clones were sequenced but 29 of those resulted in poor quality sequences or the sequenced insert was too short to give any BLAST results and were therefore discarded. Of the remaining 194 clones, 138 were up- and 56 down-regulated.

UniProt analysis of the cDNA sequences was carried out in order to identify their putative functions. Sequences without significant homology in the UniProt database were further analysed with BLASTx and BLASTn to predict their putative functions. Details of putative gene identities for the sequenced clones from the forward and reverse library are summarised in Table 2. Sequencing results for the 194 subtracted clones revealed only 3 clones (1.5%) that showed no similarities with known sequences within the UniProt and NCBI databases, and 6% corresponded to hypothetical proteins. Some of the clones were redundant and in these cases only one representative is shown. Redundancy resulted from the use of *RsaI*-restricted cDNA fragments in the SSH procedure. Because of the restriction step, two or

more different cloned cDNA fragments can represent a single transcript.

The clones that had significant sequence homologies with known sequences in the UniProt and NCBI databases were divided into seven categories through the use of the MIPS functional catalogue database (FunCatDB, Ruepp et al. 2004) and extensive search of scientific literature. The distribution of the functional classification for genes that were both up- and down-regulated is represented in Fig. 2. The largest number of induced genes was classified under the category of defence response (24%), followed by genes involved in phenylpropanoid and lignin biosynthesis (11%), and in metabolism (10%), whereas the largest number of repressed genes was classified under photosynthesis and energy (12.5%) followed by those involved in metabolism (9%). Induced genes involved in photosynthesis and energy and transport constituted the smallest group, comprising between 1 and 3% of the genes, respectively, while those genes involved in response to stimulus (8%) and those genes encoding hypothetical proteins (7%), were more abundant.

Table 2 Identification of *P. viticola*-responsive genes isolated from the SSH library in grapevine after glass slide microarray screening

Clone name	Putative function and (accession number of original matching sequence)	Gene identifiers ^b	Up/Down ^a	E-value	N ^o of clones
Defense Response					
Fw1-C9	Beta 1-3 glucanase (Q9M3U4)	XM_002277475.1	+	9.00E-67	6
Fw2-A7	Pathogenesis-related protein 10 (Q2I305)	XM_002274447	+	3.00E-57	21
Fw3-E10	Putative pathogenesis related protein 1 (Q7XAJ6)	XM_002274239	+	1.74E-66	10
Fw6-B8	Chitinase III (Q84S31)	XM_002264708	+	7.00E-96	3
Fw9-F7	F23N19.16 (Plant lipid transfer protein/Par allergen) (Q1SBU5)	XM_002264045	+	5.00E-20	5
Fw6-C6	Pleiotropic drug resistance protein 1 (NpPDR1) (Q949G3)	XM_002285142	+	4.12E-59	1
Fw11-B2	NtPRp27 (Q9XIY9) (PR-17)	XM_002269434	+	3.00E-70	2
Fw13-A12	Thaumatococcus-like protein (Q7XAU7)	XM_002283046	+	4.00E-80	1
Rv8-F12	Germin-like protein 6 (Q0PWM4)	XM_002279223	-	1.09E-50	1
Metabolism					
Fw1-D6	26 S proteasome regulatory particle triple-A ATPase subunit2b (Q8W422)	XM_002263298	+	6.00E-68	1
Fw11-H8	Maturase (Q6TNE4)	XM_002283754	+	0.71	1
Fw2-H3	Ubiquitin-conjugating enzyme (Q2I309)	XM_002281744	+	1.15E-57	2
Fw3-E5	40 S ribosomal protein S17 (Q6RJV3)	XM_002268961	+	3.00E-44	1
Fw1-E2	Monooxygenase (Monooxygenase 2) (O81816)	XM_002272572	+	4.27E-49	1
Fw5-G1	Highly similar to developmental protein DG1118 (Q9SSM4)	XM_002281246	+	6.85E-21	2
Fw6-C12	60 S ribosomal protein L13a-4 (R13A4)	XM_002268865	+	2.03E-15	2
Fw6-C8	Putative ribosomal protein S10 (Fragment) (Q0VJB2)	XM_002273250	+	2.72E-09	1
Fw5-G3	Aldehyde dehydrogenase (NAD+) (EC 1.2.1.3) (P93344)	XM_002263443	+	6.00E-57	4
Fw12-G5	Amine oxidase (Q9SW88)	XM_002263313	+	2.00E-109	5
Rv6-G2	Putative ribosomal protein S14 (Q6H7T1)	XM_002276598	-	4.97E-62	3
Rv7-F2	Putative tumor differentially expressed protein 1 (Q6K688)	XM_002270574	-	7.00E-48	2
Rv11-C12	Fiber protein Fb11 (Q8GT82)	XM_002265077	-	3.00E-24	2
Rv14-E1	IMP dehydrogenase/GMP reductase (Q1RSI8)	XM_002274242	-	7.00E-47	1
Rv7-E12	Galactinol synthase, isoform GolS-1 (Q9XGN4)	XM_002265911	-	3.80E-41	1
Rv3-E1	Cyclin-like F-box; Galactose-binding like (Q1S0J2)	XM_002279550	-	9.10E-35	2
Rv9-E3	High mobility group protein (Q4ZH67)	XM_002280048	-	1.13E-29	1
Rv14-E4	Aldo/keto reductase AKR (Q52QX9)	XM_002265991	-	5.00E-87	1
Rv9-G6	DnaJ-like protein (P93499)	XM_002265808	-	6.15E-30	1
Rv2-F6	60 S ribosomal protein L8-3 (Q42064)	XM_002269373	-	7.02E-58	1
Rv11-B2	Putative ankyrin-repeat protein (Q6TKQ6)	XM_002283462	-	1.06E-101	1
Rv5-D6	Hydrolase, alpha/beta fold family protein (Q10QA5)	XM_002285272	-	3.73E-20	1
Photosynthesis and Energy					
Fw7-F12	Oxygen-evolving enhancer protein 2, chloroplast precursor (OEE2) (Q9SLQ8)	XM_002283012	+	1.09E-100	1
Fw11-D2	Photoreceptor-interacting protein-like (Q9FJY3)	XM_002272516	+	4.00E-82	1
Fw11-D6	Chlorophyll A-B binding protein (CAB) (Q1EPK5)	XM_002274906	-	1.11E-77	10
Fw13-C7	Plastid ribosomal protein L19, putative (Q2A9H1)	XM_002265826	-	8E-18	4
Rv6-A5	Photosystem II 10 kDa polypeptide, chloroplast precursor (PII10) (Q40519)	XM_002271755	-	6.74E-45	2
Rv1-G1	Hypothetical chloroplast RF2 (Q0ZIV6)	DQ424856	-	1.44E-80	1
Rv10-F10	Type 26 kD CP29 polypeptide (Q00321)	XM_002264259	-	2.00E-47	2

Table 2 (continued)

Clone name	Putative function and (accession number of original matching sequence)	Gene identifiers ^b	Up/Down ^a	E-value	N ^o of clones
Rv4-H8	Rubisco activase (O82428)	XM_002282200	–	2.38E-90	3
Rv8-C7	Ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit (Q2I314)	XM_002276955	–	1.92E-47	2
Response to stimulus					
Fw11-A4	Methionine sulfoxide reductase A (Q6QPJ5)	XM_002278222	+	3.33E-52	1
Fw7-F4	Cysteine protease (O50002)	XM_002278588	+	3.30E-90	2
Fw1-E4	GTPase activating protein-like (Q9LVH4)	XM_002275199	+	1.46E-60	1
Fw8-B5	AtRab18 (Putative GTP-binding protein) (O23657)	XM_002267351	+	1.14E-41	1
Fw12-B9	Harpin-induced protein 1 (Q10NQ5)	XM_002265754	+	3.00E-13	1
Fw12-C9	Heavy-metal-associated domain-containing protein (AT5g14910) (Q93VK7) (HMAP)	XM_002275240	+	2.00E-13	3
Fw2-G10	Glutathione S-transferase GST (Q948X4)	XM_002262806	+	2.75E-06	2
Fw13-H6	Metallothionein-like protein 1 (MT-1)	XM_002283754	+	3.00E-04	5
Rv10-G3	AUX/IAA protein (Q1SP83)	XM_002284246	–	9.00E-70	1
Rv7-D1	Aminomethyltransferase, mitochondrial precursor (P49363)	XM_002277870	–	3.55E-71	1
Rv9-E4	Metallothionein-like protein type 3 (MT-3) (Q40256)	XM_002284942	–	1.83E-26	1
Rv8-A6	Putative metallothionein-like protein (Q9M4H3)	XM_002284942	–	2.89E-35	1
Rv2-E11	Metallothionin 2 (Q1W3C8)	XM_002265022	–	6.09E-17	1
Rv8-F5	Thioredoxin domain 2 (Q1T0K3)	XM_002266314	–	1.01E-32	1
Rv11-A12	1-aminocyclopropane-1-carboxylic acid oxidase 1 (Q84X67)	XM_002273394	–	5.41E-18	1
Rv10-B3	AOBP (Ascorbate oxidase promoter-binding protein) (Q39540)	XM_002281958	–	7.92E-39	1
Phenylpropanoid and Lignin Biosynthesis					
Fw1-F6	Chalcone synthase (O80407)	XM_002276910	+	1.00E-98	7
Fw7-B2	Pectin methylesterase inhibitor isoform (Q0Z7S4)	XM_002264168	+	1.15E-12	1
Fw2-D8	Expansin 45, endoglucanase-like (Q1RYX8)	XM_002284960	+	3.03E-13	1
Fw2-E10	Cinnamate-4-hydroxylase (Q3HM04)	XM_002266202	+	9.00E-106	3
Fw3-C10	Naringenin,2-oxoglutarate 3-dioxygenase (Flavonone-3-hydroxylase)	XM_002267604	+	1.00E-90	2
Fw9-E1	Putative pectin methylesterase (Q8H223)	XR_077457	+	3.00E-06	1
Fw11-G10	Chalcone isomerase	XM_002282072	+	2.00E-70	7
Signal Perception and Transduction					
Fw5-G7	Expressed protein (Leucine Rich Repeat family protein) (Q53PD8)	XM_002268472	+	2.36E-06	1
Fw5-F6	Sucrose responsive element binding protein (Q1XAN1)	XM_002283311	+	8.66E-63	1
Fw2-F11	Putative receptor-like protein kinase 1 (Q5JN27)	XM_002266170	+	9.19E-68	1
Fw2-A10	Putative peroxisomal membrane protein(22-kDa)(PMP22) (Q6Z1N7)	XM_002269300	+	3.00E-47	4
Fw11-A10	Membrane protein (Q2M5F0)	XM_002277120	+	2.90E-19	1
Fw14-B7	Zinc finger, C2H2 type family protein, expressed (Q2QX40)	AM482928	+	3.3	1
Fw8-E7	SOS2-like protein kinase (Q8LK24)	XM_002285460	+	5.00E-24	1
Fw1-C1	Cyclin-like F-box; Serine/threonine protein phosphatase, BSU1 (Q1RZ31)	XM_002265077	+	1.19E-49	1
Rv12-G10	Phosphate translocator precursor (Q40568)	XM_002278793	–	3.00E-56	1
Transport					
Fw5-C10	Sorbitol transporter (Q84KI7)	XM_002284527	+	4.29E-14	1
Fw2-B9	Putative peptide transporter protein (Q8VZN7)	XM_002266869	+	4.22E-20	2
Fw11-D5	Copper transporter 1 (Q547P9)	XM_002280052	+	4.00E-17	2

Table 2 (continued)

Clone name	Putative function and (accession number of original matching sequence)	Gene identifiers ^b	Up/Down ^a	E-value	N ^o of clones
Unclassified					
Fw9-C9	Hypothetical protein T6H20.190 (Q9STF2)	XM_002282694	+	1.10E-38	1
Fw2-E9	T17H3.3 protein (Hypothetical protein) (Similar to CGI-126 protein) (Q9SXC8)	XM_002285545	+	3.00E-29	1
Fw6-D5	Hypothetical protein (At3g15810) (Q9LVZ8)	XM_002266546	+	1.00E-22	1
Fw6-E1	Hypothetical protein T16K5.140 (Hypothetical protein At3g49790) (Q9M2X9)	AM438165	+	2.2	1
Fw9-G4	Hypothetical protein rps12 (Q3BAI4)	AM483961	+	3.00E-27	1
Fw9-G9	Hypothetical protein At2g39140 (O80967)	XM_002270150	+	1.02E-24	1
Fw4-H6	Hypothetical protein (Q56ZH1)	XM_002275456	+	1.24E-06	1
Fw10-D4	Hypothetical protein (Q3LSN3)	XM_002279821	+	3.00E-60	1
Fw2-C7	Unnamed protein product (CAO38708.1)	XM_002265072	+	4.00E-19	1
Fw1-F10	Hypothetical protein ORF1 (Q9FSX3)	XM_002263298	+	7.00E-18	1
Fw1-F2	No homology		+		1
Fw2-C10	No homology		+		1
Fw1-E6	No homology		+		1
Rv7-H8	Expressed protein (Q10Q41)	XM_002271976	-	2.77E-28	1
Rv3-E12	Hypothetical protein (Q8LDM9)	XM_002283263	-	2.54E-07	1
Rv8-D10	Hypothetical protein (Q1SFB2)	XM_002270765	-	1.45E-10	1
Rv4-A11	Hypothetical protein (CAN65189)	XM_002266869	-	4.00E-94	1
Rv9-G5	No homology		-		1

^a“+” means gene was induced by *P. viticola* from screening of the SSH libraries, and comes from the forward library, “-” means gene was suppressed by *P. viticola* from screening of the SSH libraries and comes from the reverse library

^bGrapevine gene identifiers were annotated according to the NCBI database

Expression analysis of differentially responsive genes to *P. viticola* infection by RT-qPCR

To analyse the expression patterns of a number of genes during the interaction of *V. vinifera* with *P. viticola*, reverse transcriptase quantitative real-time PCR (RT-qPCR) was performed on inoculated and mock-inoculated samples 0, 8, 12, 24 and 48 h after *P. viticola* inoculation. The selected genes corresponded to a range of functional categories: defence response (6 genes), phenylpropanoid and lignin biosynthesis (4 genes) and response to stimulus (2 genes) (Fig. 3). RT-qPCR was carried out with two biological replicates. Total RNA extracted from grapevine seedlings was reverse-transcribed into first-strand cDNA, and used as template for quantitative RT-PCR analysis. Results of the time-course experiment showed that PR-genes were highly induced compared to those involved in phenylpropanoid and lignin biosynthesis and in response to stimulus, which was

in good agreement with the library verification results (data not shown). Among the PR-genes, a β -1,3-glucanase (*Gluc*) and a pathogenesis-related protein 10 (*PR-10*) showed the highest fold increase in expression level. The expression level of *Gluc* increased transiently over the 48 h time course, whereas the expression level of *PR-10* reached a maximum level at 12 hpi, decreasing at 24 hpi and remaining constant up to 48 h. The expression pattern of a chitinase III (*Chit III*) was similar to that of *PR-10* with the exception that after a slight decrease in expression level at 24 hpi, it increased gradually until 48 hpi. Another PR gene, pathogenesis-related protein 1 (*PR-1*) was induced 9.5-fold at 12 hpi where it reached maximum level and then decreased reaching levels close to the non-infected samples. Like *Gluc* the expression level of a pathogenesis-related protein 17 (*PR-17*) was transiently induced over the 48 h time course. The only defence-related gene which did not show any significant change in expression level over




Fig. 2 Functional classification of genes up- and down-regulated in grapevine by *P. viticola* inoculation. Differentially expressed genes were classified into seven groups and the percentage of the genes within each category is shown

the time course was a thaumatin-like protein (*PR-5*). Similarly, one of the four genes involved in phenylpropanoid and lignin biosynthesis, namely a chalcone synthase (*CHS*) was down-regulated. Flavonone-3-hydroxylase (*F3H*) and cinnamate-4-hydroxylase (*C4H*) were not highly induced over the 48 h time course, both showed a 2-fold increase in expression level at 24 hpi, whereas chalcone isomerase (*CHI*) was induced 8-fold at 24 hpi. Of the two genes involved in response to stimulus, only a metallothionein-like protein (*MT*) showed a 2-fold increase in expression at 12 hpi after which the expression level decreased reaching levels of the non-infected samples at 48 hpi, whereas a heavy-metal-associated protein (*HMAP*) showed no significant change in expression level between infected and non-infected plants.

Microscopic observation of the response of the susceptible grapevine cv. Chasselas to *P. viticola* infection

To monitor response mechanisms and pathogen growth in leaves of a susceptible grapevine, leaf disks were sampled at different time points post inoculation. The development of *P. viticola* in the susceptible grapevine

caused visible symptoms such as curling of a heavy infected leaf, oil spots which started to undergo necrosis and sporulation on the abaxial leaf surface within 7 days after inoculation (Fig. 4a, b & c). After 4 dpi hyphal growth was observed (Fig. 4d) and after 7 dpi extensive colonization of leaf tissues was observed by the presence of abundant branched hyphae and the formation of sporangiophores bearing a large number of sporangia (Fig. 4e & g). Aggregation of hyphae in the sub-stomatal cavity was observed (Fig. 4f). As expected mock-inoculated samples at 7 dpi showed no signs of infection (Fig. 4o).

Lignification of the leaf tissues was observed by phloroglucinol-HCl staining leading to a red colourization of lignin. In the inoculated samples 2 and 4 dpi very few red-stained cells were observed (Fig. 4h & i). However at 7 dpi prominent red-stained cells were observed (Fig. 4j) evenly spread throughout the host cells and in close proximity to a stoma (Fig. 4k). No lignification was observed in the mock-inoculated samples at 7 dpi (Fig. 4p).


Plant tissues stained with Naturstoff display a light yellow fluorescence that shows the presence of flavonoids. As expected, no accumulation of flavonoids was found in the mock-inoculated samples 7dpi (Fig. 4q) nor in the inoculated samples 2 dpi (Fig. 4l) but at 4 dpi the presence of the light yellow fluorescence could be seen in the inoculated samples (Fig. 4m) and by 7 dpi a predominant increase in the light yellow fluorescence could be observed (Fig. 4n).

Discussion

Compatibility describes the complementary relationship between a plant species and an adapted pathogen which ultimately results in disease. The progression of *P. viticola* culminating in the appearance of oil spots (Fig. 4b) and the formation of fully developed sporangiophores with sporangia (Fig. 4e) in *V. vinifera* cv. Chasselas represents a compatible interaction. Despite considerable recent progress, our understanding of many aspects of plant-birotroph compatibility is still in its early stages.

Obligate biotrophs, such as powdery and downy mildews and rust fungi, cannot be extensively cultured *in vitro*, since they have to form specialized infection structures, haustoria, within infected host cells. In contrast to necrotrophic and hemibiotrophic fungal

Fig. 3 Reverse transcriptase quantitative real-time PCR (RT-qPCR) analysis of transcript accumulation in grapevine in response to *P. viticola* challenge. Transcript accumulation of *Gluc*, *PR-10*, *Chit III*, *PR-1*, *PR-17*, *PR-5*, *CHS*, *C4H*, *CHI*, *F3H*, *MT* and *HMAP* genes was monitored in mock-inoculated (dashed lines) and *P. viticola*-inoculated (solid line) plants. Transcript levels of the twelve genes are expressed as relative values normalized to the transcript level of *EF1- α* gene, used as an internal reference. Results are means of triplicate data from one representative experiment among two independent repetitions



pathogens, obligate biotrophs are entirely dependent on living plant tissue for their growth and propagation. This lack of host cell death in plant-biophyte interactions might be accomplished by the pathogen's ability to avoid or durably suppress pre-formed and induced host

defences (Mendgen and Hahn 2002). It is therefore probable that specific host genes and/or proteins are targeted by biotrophs to achieve these goals, and so be considered as compatibility factors that are essential for successful pathogenesis (Panstruga 2003).

Construction and screening of a SSH library at 24 h after *P. viticola* inoculation

Gene expression profiling has become a valuable tool in functional genomics. SSH is a powerful method to isolate genes that are specifically expressed under diverse sets of conditions. To better understand the molecular basis of compatibility and disease development in a grapevine-*Plasmopara* interaction, we used SSH to identify cDNAs that were differentially expressed in infected plants 24 h after inoculation. Recently, it was shown that rice defence genes induced in resistant plants were also induced in susceptible interactions (Vergne et al. 2007). Moreover, Jantasuriyarat et al. (2005) show that the transcriptional response of a susceptible plant at 24 hpi is most similar to the response of a resistant plant at 6 hpi. They also found that the most highly induced rice genes at 24 hpi (50-fold induction) in a susceptible interaction are also induced in resistant plants (6 and 24 hpi).

Downy mildew oomycetes enter the intercellular spaces of the leaf mesophyll via stomata where they form an infection vesicle with primary hypha and the first haustorium. This mode of entry is expected to have consequences on the timing of recognition of the pathogen by the plant cells and on the nature of induced defence responses. The outcome of plant-pathogen interaction seems to be determined at the time of the subsequent penetration of intact host cells by the haustorium. Recently, two microscopy studies of the infection process of *P. viticola* showed that elongated hyphae with haustoria invaded the intercellular space of the mesophyll by 24 hpi in the resistant *Vitis riparia* cv. Gloire de Montpellier and in the susceptible *V. vinifera* cv. Muscat Ottonel (Diez-Navajas et al. 2008; Unger et al. 2007). Since it is known that the first infection steps, occurring within 24 h after inoculation are essentially the same in a resistant and in a susceptible cultivar we decided to construct the SSH library at 24 hpi. Microscopically, the main difference in infection between the two mentioned cultivars occurs within 48–72 hpi.

Recently, Polesani et al. (2008) used a cDNA-AFLP method to investigate the compatible interaction between *P. viticola* and grapevine at the oil spot stage. They found that nearly 70% of the identified differentially expressed grapevine genes were down-regulated during infection due to the exploitation of

cellular resources and the suppression of defence responses. We only found 30% and 13% of the differentially expressed genes to be down-regulated in the forward and reverse library, respectively, the main difference between these two studies being that Polesani and associates carried out their analyses at the oil spot stage, where infection has already been established, whereas our analyses was done at 24 h after inoculation, which mirrors the early stages of infection. At this time point we still expect the susceptible cultivar to try to mount a defence response.

Since the majority of the methods used to screen a SSH library involves visual inspection of inverse dot blots, this constitutes a potential source of false positives. We chose a quantitative screening method that was developed to evaluate the quality of the SSH libraries (Van den Berg et al. 2004; Berger et al. 2007). This approach facilitates the choice of clones from the library for further analysis, such as DNA sequencing, RT-qPCR, Northern blotting or detailed expression profiling using a custom cDNA microarray (Berger et al. 2007). Another advantage of using this screening method with the software SSHscreen is the inclusion of statistical tests of the microarray data to provide further confidence in the choice of genes.

Defence-related genes

The differentially expressed cDNAs isolated from the SSH were classified into several categories and were involved in a number of physiological and molecular events including: metabolism, photosynthesis and energy, response to stimulus, phenylpropanoid and lignin biosynthesis, signal perception and transduction, transport and defence response. A significant number of genes identified in this study that were involved in defence included PR proteins (PR-1, PR-5, PR-10 and PR-17). PR proteins were identified several years ago as being associated with resistance of plants to various pathogens and abiotic stresses (van Loon et al. 2006). Members of the PR-1, and the thaumatin-like PR-5 families have been associated with activity against oomycetes. PR-5-type proteins are thought to create trans-membrane pores and exhibit antifungal activities *in vitro* by blocking spore germination and germ tube growth of *Uncinula necator*, *Phomopsis viticola* and *Botrytis cinerea* (Monteiro et al. 2003). Interestingly, we found *PR-1*


to be induced as early as 12 hpi whereas *PR-5* showed no induction over the 48 h time course after inoculation with *P. viticola*. The latter is in agreement with Kortekamp (2006), who found no difference in *PR-5* expression in a susceptible and resistant cultivar after inoculation with *P. viticola* indicating that the *PR-5*'s role in defence against *P. viticola* remains unclear. The role of *PR-1* in defence against *P. viticola* also remains uncertain. Interestingly, *P. viticola* did not induce the expression of *PR-1* in the susceptible cultivar Chardonnay according to Chong et al. (2008) but *PR-1* was detected 24 h following inoculation with *P. viticola* in *V. vinifera* cv. Riesling (Wielgoss and Kortekamp 2006). These differences in *PR-1* expression may be related to the use of different cultivars, different isolates of *P. viticola*, semi-quantitative versus quantitative RT-PCR, or even different time points after inoculation. Chong and associates looked at *PR-1* expression between 18 and 96 hpi and we found that at 48 hpi the expression level of *PR-1* was already returning to basal levels close to those of the non-inoculated plants.

A large number of *PR-10*-related sequences have been identified in many plants. Sequence comparison indicated that *PR-10* proteins are structurally related to ribonucleases. The inducible expression of *PR-10* genes in response to attacks by pathogens has been widely investigated in a number of plant species. Pathogens triggering a *PR-10* response include viruses, bacteria and fungi (Liu and Ekramoddoullah 2006). Some *PR-10* proteins display constitutive expression patterns unrelated to the pathogen response. It has been shown that *PR-10* proteins are also developmentally regulated in different plant tissues and organs, therefore it is not reasonable to assign a unique function to *PR-10* proteins throughout the plant kingdom. Nevertheless, *PR-10* expression was induced in grapevine after challenge with *P. viticola*. A strong induction of *PR-10* was observed at 12 hpi which is in agreement with a number of other studies that found transcript accumulation of *PR-10* in the susceptible grapevine cv. Riesling 12 hpi with *P. viticola* (Kortekamp 2006), in rice within 12 hpi with *Magnaporthe grisea* (McGee et al. 2001) and in sorghum, inducible *PR-10* transcripts reached a maximum level at 12 hpi (Lo et al. 1999). When taking these results into account, the induction of *PR-10* may play an important role in the defence of grapevine against *P. viticola*.

Fig. 4 Development of *P. viticola* and accumulation of lignin and flavonoids in the susceptible cultivar Chasselas. Plants were inoculated with sporangia of *P. viticola* (4×10^4 sporangia ml^{-1}) (a–n), sampled 2 dpi (h and l), 4 dpi (d, i and m) and 7 dpi (a–c, e, f, g, i, j, k, and n). Mock-inoculated plants were sampled at 7 d (o–q). Infected leaves were stained with lactophenol-trypan blue and analyzed by light microscopy (d–g), stained with phloroglucinol-HCL and analyzed by light microscopy (h–k and p), or stained with Naturstoff reagent A and analyzed by epifluorescence microscopy (l–n and q). a White sporulation of *P. viticola* on the abaxial leaf surface; b heavily infected leaf showing typical curling symptoms; c oil spots starting to undergo necrosis on the adaxial leaf surface. d Initial hyphal growth, e extensive colonization of leaf tissues and abundant branched hyphae, f aggregation of hyphae (hy) in the sub-stomatal cavity (st), g sporangiophores bearing a large number of sporangia. h Very little amount of lignin, i presence of small amount of lignin as shown by red-coloured tissue, j increased amount of red-stained lignin in tissues, k prominent lignin deposition close to a stoma. l No presence of flavonoids at 2 dpi, m presence of flavonoids as indicated by yellow fluorescence (4 dpi), n accumulation of yellow fluorescence. o No presence of infection in mock-inoculated plants at 7d, p and q no presence of lignin and flavonoids in mock-inoculated plants respectively

Many pathogenesis-related proteins were first characterised in tobacco, NtPRp27, is constitutively expressed in tobacco roots but can be induced by tobacco mosaic virus, wounding, drought and by the application of ethylene, methyl jasmonate, salicylic acid and abscisic acid (Okushima et al. 2000). Recently NtPRp27 became the prototype for the new PR family PR-17. Accumulation of mRNA encoding PR-17 proteins has been reported after benzothiadiazole treatment, inoculation with *Erysiphe graminis* f. sp. *tritici* in wheat (Gorlach et al. 1996) and inoculation with *Blumeria graminis* f. sp. *hordei* (*Bgh*) in barley (Christensen et al. 2002). The transient accumulation of *PR-17* after *P. viticola* inoculation was in close agreement with both transcript and protein accumulation in the compatible and incompatible interactions in barley in response to *Bgh*, diversions in these interactions occurred after 72 hpi.

Two other PR proteins namely a *chitinase III* (*PR-8*) and a β -1,3-*glucanase* (*PR-2*) were induced after *P. viticola* inoculation. Both PR proteins are thought to provide defence against fungal pathogens by hydrolysing components of the cell walls of invading fungal hyphae (Mauch et al. 1988). It has been shown that cell walls of *P. viticola* consist of chitin and β -1,3-*glucan* as seen on the basis of histochemical staining with specific fluorochromes, (Kortekamp 2005) therefore expression of both *chitinase III* and



β -1,3-glucanase may inhibit fungal growth. The induction of β -1,3-glucanase was transient and to much higher levels than that found for *chitinase III*. Interestingly, Kortekamp (2005) showed that *P. viticola* septa are composed of β -1,3-glucans but not chitin, which could trigger the host plant to increase the production of β -1,3-glucanase. Our results for *chitinase III* are in accordance with those of Busam et al. (1997), showing that its induction by *P. viticola* in both a susceptible and resistant cultivar. They also showed that *chitinase III* increased transiently in the healthy tissue of younger, next-stage leaves and that *chitinase III* was up-regulated in response to treatment with INA and BTH. Overall the results suggest that the selective expression of *chitinase III* is a reliable indicator of the systemic acquired resistance (SAR) response in grapevine.

We identified two other homologs involved in defence, namely a lipid transfer protein (LTP) and a pleiotropic drug resistance protein 1 (NpPDR1). LTPs are members of the family of pathogenesis-related proteins (PR-14) that are involved in plant defence responses. Plant LTPs also have developmental and tissue specificities, they are modulated by environmental and stress conditions and have a number of diverse functions, including cutin biosynthesis, surface wax formation and adaptation of the plant to environmental changes. An interesting feature of LTPs is their structural similarity with elicitor, which is a small lipid-binding protein that is secreted by the phytopathogenic oomycetes *Phytophthora* and *Pythium*, and which triggers the hypersensitive response (HR) and SAR in tobacco (Ponchet et al. 1999). A homologous gene encoding a protein of the pleiotropic drug resistance (PDR) subfamily of ATP-binding cassette (ABC) transporters was found to be induced in grapevine by *P. viticola*. These ABC transporters mediate the translocation of a diverse array of compounds across biological membranes, including toxins, drugs, glutathione conjugates, peptides and secondary metabolites (Yazaki 2006). There is also increasing evidence for the involvement of ABC transporters in defence. Stukkens et al. (2005) found that a PDR protein in tobacco leaves (NpPDR1) was induced following pathogen attack and silencing of NpPDR1 by RNA interference resulted in increased sensitivity to the plant pathogen *Botrytis cinerea*. More recently Kobae et al. (2006) demonstrated that a loss of an *Arabidopsis* plasma membrane-localized PDR ABC

transporter (AtPDR8) lead to hypersensitive cell death upon pathogen infection.

The only defence-related gene we found to be down-regulated in response to *P. viticola* infection was a germin-like protein 6 (GLP). In grapevine seven members of the GLP gene family (*VvGLP*) have been identified (Godfrey et al. 2007). *GLP* genes are expressed in various organs and developmental stages in plants and in response to a number of biotic and abiotic stresses (Bernier and Berna 2001). In grapevine, expression analysis revealed that the *VvGLP* genes exhibited diverse and highly specific patterns of expression in response to a variety of abiotic and biotic treatments, including challenge by *Erysiphe necator*, *P. viticola* and *B. cinerea*. Down regulation of *VvGLP6* in leaf tissue infected with *P. viticola* has been reported previously (Godfrey et al. 2007).

Genes involved in the phenylpropanoid and lignin biosynthesis pathway

In the general phenylpropanoid pathway, three enzymatic transformations redirect the carbon flow from primary metabolism, transforming phenylalanine into the coenzyme A (CoA)-activated hydroxycinnamoyl thioester capable of entering the two major downstream pathways: monolignol and flavonoid biosynthesis. The pathway that provides the lignin-building monolignol units is strongly activated after infection by pathogens or treatments with elicitors. In infected plants, deposition of phenylpropanoid compounds is a part of the cell wall reinforcement that restricts pathogen invasion. Accumulation of soluble and cell-wall bound phenolics in plant tissues challenged by fungal pathogens has been suggested as a defence strategy. Enzymes such as phenylalanine ammonia lyase (PAL; EC 4.3.1.5), cinnamate-4-hydroxylase (C4H; EC 1.14.13.11) and 4-coumarate:coenzyme A ligase (4CL, EC 6.2.1.12) are crucial to phenylpropanoid metabolism. We found the expression of *C4H* to be induced upon *P. viticola* challenge peaking at 12 hpi. Sewalt et al. (1997) showed that reduced C4H activity correlated with reduced levels of lignin indicating that there could be a correlation between *C4H* expression and lignification. Along with the induced expression of *C4H* we found an increase in lignin deposition as shown through histochemical staining (Fig. 4h–k).

All flavonoids are derived from the chalcone scaffold, which is biosynthesized by the enzyme chalcone synthase (CHS). Following CHS, an assemblage of isomerases (CHI), reductases, hydroxylases (F3H), glycosyltransferases and acyltransferases make up the basic flavonoid skeleton leading to a number of flavonoid chemical subclasses such as flavones, isoflavanoids, flavonols, flavandiols, proanthocyanidins and anthocyanins. Even though we found the expression of *CHS* to be down-regulated, the other two enzymes which are involved down-stream from *CHS*, namely *CHI* and *F3H*, were induced by *P. viticola* challenge. The expression level of both *CHI* and *F3H* peaked at 24 hpi. The expression of these two genes coincided with the production of flavonoids as seen by histochemical staining (Fig. 4l–n).

In grapevine, among the best characterised defence reactions upon fungal infection, is the accumulation of stilbene phytoalexins. Resistant grape cultivars artificially inoculated with *P. viticola* showed high amounts of stilbenic phytoalexins at the site of infection, and these phytoalexins have significant inhibitory effects on the mobility of *P. viticola* zoospores (Pezet et al. 2004). Chalcone synthase and stilbene synthase (*STS*) are further key branch-point enzymes in the phenylpropanoid pathway and their enzymatic activities and gene expression are related to phytoalexin biosynthesis in grapevine. We expect to find *STS* present in the SSH library since in separate studies, expression of *STS* in grapevine was induced after *P. viticola* infection (Chong et al. 2008; Kortekamp 2006). We have consistently seen that the expression of *STS* in *P. viticola*-inoculated grapevine seedlings decreases drastically at 24 hpi and remains low up to 48 hpi (Slaughter et al., unpublished results). Interestingly, Chong et al. (2008) found the expression of *STS* to be induced 48 h after *P. viticola* inoculation and not before and for that reason we were not able to identify *STS* in our library. We found the expression of *CHS* to be down-regulated and we did not find the presence of *STS* and since both are key enzymes in the production of stilbene phytoalexins, they could play an important role in the resistance of grapevine to *P. viticola* but since the SSH library was constructed from a susceptible cultivar to *P. viticola*, the down-regulation of *CHS* and the decrease in *STS* expression could contribute to the susceptibility of this cultivar to *P. viticola*.

Response to stimulus-related genes

Plant metallothioneins (MTs) are small, highly conserved cysteine-rich heavy-metal-binding proteins, typically classified into four categories (types 1 to 4). Plant MT gene expression is regulated by various factors, including metal ions, developmental stages, symbiotic interaction, and various stress responses. Stresses such as wounding and virus infection were shown to induce the expression of MT in tobacco, while differential expression of MT was detected in *Arabidopsis* following infection of compatible and incompatible strains of *Peronospora* spp. In this respect plant MTs have been proposed to function as reactive oxygen species (ROS) scavengers that protect plant DNA from oxidative damage caused by free radicals released during the oxidative burst (Wong et al. 2004). Although the majority of expressed MT genes belong to type 2, type 1, 3 and 4 MT have also been reported to be involved in various plant processes. Transcript accumulation of type 1 MT was shown in various organs and also in response to pathogen attack (Butt et al. 1998). In this study, induction of a type 1 MT was observed 12 hpi followed by a decline at 24 hpi. A plausible explanation is that the plant accumulated large amounts of MT transcripts in response to *P. viticola* within early stages of infection, but mRNA levels rapidly returned to levels comparable with those expressed in control plants. A similar trend of MT differential gene expression in tobacco plants following *Tobacco mosaic virus* (TMV) infection over a period of 72 h was observed (Choi et al. 1996). Relative amounts of MT mRNA were highly induced at 12 h, sharply dropped after 24 h following TMV infection and remained the same thereafter. We found the expression of a type 2 MT to be down-regulated in response to *P. viticola* challenge, which is in accordance with Wong et al. (2004) who found in rice the expression of *OsMT2b* to be synergistically down-regulated by *OsRac1* and rice blast-derived elicitors. They also demonstrated increased susceptibility to bacterial blight and blast fungus in transgenic plants overexpressing *OsMT2b*. We also found a type 3 MT to be down-regulated in response to *P. viticola* challenge. So far there has only been one case in plants where a type 3 MT was inducible in response to pathogen attack, in velvetleaf leaves in response to *Colletotrichum coccodes*. (Dauch and Jabaji-Hare,

2006). Up-regulation of plant type 3 MT so far has been mostly observed as part of senescence-related responses in plants. In the case of a heavy-metal-associated domain-containing protein we did not find any induction between inoculated and control plants.

Suppression of photosynthesis-related genes

In this study, the largest group of genes suppressed in grapevine during *P. viticola* challenge were photosynthesis-related, namely chlorophyll a-b binding protein, photosystem II polypeptide and the CP29 subunit of photosystem II. Transcriptional down-regulation of photosynthesis-related genes has previously been reported for incompatible and compatible interactions (Moy et al. 2004; Restrepo et al. 2005; Vergne et al. 2007). A down-regulation of genes related to photosynthesis was shown by Polesani et al. (2008) in grapevine at the oil spot stage after challenge with *P. viticola*. The reason for the reduction in photosynthesis during the compatible interaction between grapevine and *P. viticola* is unknown, but Fung et al. (2008) have suggested that in a compatible interaction of grapevine with powdery mildew, there is an up-regulation of invertases, which are involved in degradation of carbon reserves into hexoses, resulting in the reduction of photosynthetic rates. This metabolic shift seems to be coordinated with up-regulation of genes involved in the synthesis of secondary metabolites (flavonoids and lignin) via the oxidative pentose phosphate and shikimate pathway. Increased invertase activity leads to carbohydrate accumulation, which in turn may inhibit the Calvin cycle, which also limits photosynthesis. Two genes encoding enzymes involved in the Calvin cycle, namely rubisco activase and ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit were down-regulated.

Metabolism-associated genes

Proteolysis of important regulatory proteins is a key aspect of cellular regulation in eukaryotes and there is evidence that the ubiquitin-proteasome pathways are important in implementation of the plant defence response. In response to pathogen attack, the Ub/26 S proteasome pathway initiates programmed cell death to localise pathogen spread (Smalle and Vierstra 2004). We found homologs of a 26 S proteasome

and an ubiquitin-conjugating enzyme to be up-regulated in grapevine in response to *P. viticola*.

In summary, the SSH method has allowed us to generate a differential cDNA library enriched for *Plasmopara*-responsive genes from grapevine seedlings. This is the first large-scale investigation into the early response (24 hpi) of compatibility between *V. vinifera* and the biotrophic pathogen *P. viticola*. Polesani et al. (2008) performed cDNA-AFLP analysis on a compatible interaction of grapevine with *P. viticola* on the oil spot stage of infection which follows 2–3 weeks after inoculation; this could be defined as a late response to pathogen challenge, whereas we were interested in analyzing the early response to *P. viticola* challenge. Even though we are dealing with a susceptible cultivar to *P. viticola* challenge, the host plant was still able to mount a defence response as seen by the number of genes identified which are involved in defence mechanisms. Further characterization and functional analysis of the genes generated from the library will facilitate our understanding of the defence mechanisms in grapevine plants to *P. viticola* infection.

Acknowledgements This project was funded by the National Centre of Competence in Research (NCCR) Plant Survival, a research programme of the Swiss National Science Foundation. We thank Mrs. M. Waldner (Syngenta, Stein, Switzerland) for the grapevine seedlings and Dr J. Weber (Lausanne DNA Array Facility, University of Lausanne) for the assistance in the making of the SSH slides for the microarray screening.

References

- Adams, M. D., Kelley, J. M., Gocayne, J. D., Dubnick, M., Polymeropoulos, M. H., Xiao, H., et al. (1991). Complementary DNA sequencing: expressed sequence tags and human genome project. *Science*, 252, 1651–1656.
- Allegre, M., Daire, X., Heloir, M. C., Trouvelot, S., Mercier, L., Adrian, M., et al. (2007). Stomatal deregulation in *Plasmopara viticola*-infected grapevine leaves. *The New Phytologist*, 173, 832–840.
- Aziz, A., Poinssot, B., Daire, X., Adrian, M., Bezier, A., Lambert, B., et al. (2003). Laminarin elicits defense responses in grapevine and induces protection against *Botrytis cinerea* and *Plasmopara viticola*. *Molecular Plant-Microbe Interactions*, 16, 1118–1128.
- Berger, D. K., Crampton, B. G., Hein, I., & Vos, W. (2007). Screening of cDNA libraries on glass slide microarrays. *Methods in Molecular Biology*, 382, 177–203.
- Bernier, F., & Berna, A. (2001). Germins and germin-like proteins: Plant do-all proteins. But what do they do exactly? *Plant Physiology and Biochemistry*, 39, 545–554.

- Birch, P. R. J., Avrova, A. O., Duncan, J. M., Lyon, G. D., & Toth, R. L. (1999). Isolation of potato genes that are induced during an early stage of the hypersensitive response to *Phytophthora infestans*. *Molecular Plant-Microbe Interactions*, *12*, 356–361.
- Brenner, S., Johnson, M., Bridgham, J., Golda, G., Lloyd, D. H., Johnson, D., et al. (2000). Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays. *Nature Biotechnology*, *18*, 630–634.
- Busam, G., Kassemeyer, H. H., & Matern, U. (1997). Differential expression of chitinases in *Vitis vinifera* L. responding to systemic acquired resistance activators or fungal challenge. *Plant Physiology*, *115*, 1029–1038.
- Butt, A., Mousley, C., Morris, K., Beynon, J., Can, C., Holub, E., et al. (1998). Differential expression of a senescence-enhanced metallothionein gene in *Arabidopsis* in response to isolates of *Peronospora parasitica* and *Pseudomonas syringae*. *The Plant Journal*, *16*, 209–221.
- Choi, D., Kim, H. M., Yun, H. K., Park, J. A., Kim, W. T., & Bok, S. H. (1996). Molecular cloning of a metallothionein-like gene from *Nicotiana glutinosa* L. and its induction by wounding and tobacco mosaic virus infection. *Plant Physiology*, *112*, 353–359.
- Chong, J., Le Henaff, G., Bertsch, C., & Walter, B. (2008). Identification, expression analysis and characterization of defense and signaling genes in *Vitis vinifera*. *Plant Physiology and Biochemistry*, *46*, 469–481.
- Christensen, A. B., Cho, B. H., Naesby, M., Gregersen, P. L., Brandt, J., Madriz-Ordenana, K., et al. (2002). The molecular characterization of two barley proteins establishes the novel PR-17 family of pathogenesis-related proteins. *Molecular Plant Pathology*, *3*, 135–144.
- Coetzer, N., Gazendam, I., Oelofse, D., & Berger, D. K. (2010). SSHscreen and SSHdb, generic software for microarray based gene discovery: application to the stress response in cowpea. *Plant Methods*, *6*, 10.
- Dai, G. H., Andary, C., Mondolot-Cosson, L., & Boubals, D. (1995). Histochemical studies on the interaction between three species of grapevine *Vitis vinifera*, *V. rupestris* and *V. rotundifolia* and the downy fungus, *Plasmopara viticola*. *Physiological and Molecular Plant Pathology*, *46*, 177–188.
- Dauch, A. L., & Jabaji-Hare, S. H. (2006). Metallothionein and bZIP Transcription Factor Genes from Velvetleaf and Their Differential Expression Following *Colletotrichum coccodes* Infection. *Phytopathology*, *96*, 1116–1123.
- Degenhardt, J., Al-Masri, A. N., Kurkcuoglu, S., Szankowski, I., & Gau, A. E. (2005). Characterization by suppression subtractive hybridization of transcripts that are differentially expressed in leaves of apple scab-resistant and susceptible cultivars of *Malus domestica*. *Molecular Genetics and Genomics*, *273*, 326–335.
- Derckel, J. P., Baillieux, F., Manteau, S., Audran, J. C., Haye, B., Lambert, B., et al. (1999). Differential Induction of Grapevine Defenses by Two Strains of *Botrytis cinerea*. *Phytopathology*, *89*, 197–203.
- Diez-Navajas, A. M., Wiedemann-Merdinoglu, S., Greif, C., & Merdinoglu, D. (2008). Nonhost versus host resistance to the grapevine downy mildew, *Plasmopara viticola*, studied at the tissue level. *Phytopathology*, *98*, 776–780.
- Fung, R. W., Gonzalo, M., Fekete, C., Kovacs, L. G., He, Y., Marsh, E., et al. (2008). Powdery mildew induces defense-oriented reprogramming of the transcriptome in a susceptible but not in a resistant grapevine. *Plant Physiology*, *146*, 236–249.
- Gindro, K., Pezet, R., & Viret, O. (2003). Histological study of the responses of two *Vitis vinifera* cultivars (resistant and susceptible) to *Plasmopara viticola* infections. *Plant Physiology and Biochemistry*, *41*, 846–853.
- Glazebrook, J. (2005). Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annual Review of Phytopathology*, *43*, 205–227.
- Godfrey, D., Able, A. J., & Dry, I. B. (2007). Induction of a grapevine germin-like protein (VvGLP3) gene is closely linked to the site of *Erysiphe necator* infection: a possible role in defense? *Molecular Plant-Microbe Interactions*, *20*, 1112–1125.
- Gorlach, J., Volrath, S., Knauf-Beiter, G., Hengy, G., Beckhove, U., Kogel, K. H., et al. (1996). Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *The Plant Cell*, *8*, 629–643.
- Hamiduzzaman, M. M., Jakab, G., Barnavon, L., Neuhaus, J. M., & Mauch-Mani, B. (2005). beta-Aminobutyric acid-induced resistance against downy mildew in grapevine acts through the potentiation of callose formation and jasmonic acid signaling. *Molecular Plant-Microbe Interactions*, *18*, 819–829.
- Heath, M. C. (2000). Nonhost resistance and nonspecific plant defenses. *Current Opinion in Plant Biology*, *3*, 315–319.
- Iandolino, A. B., Goes da Silva, F., Lim, H., Choi, H., Williams, L. E., & Cook, D. R. (2004). High-quality RNA, cDNA, and derived EST libraries from grapevine (*Vitis vinifera* L.). *Plant Molecular Biology Reporter*, *22*, 269–278.
- Jantasuriyarat, C., Gowda, M., Haller, K., Hatfield, J., Lu, G., Stahlberg, E., et al. (2005). Large-scale identification of expressed sequence tags involved in rice and rice blast fungus interaction. *Plant Physiology*, *138*, 105–115.
- Keogh, R. C., Deverall, B. J., & McLeod, S. (1980). Comparison of histological and physiological responses to *Phakopsora pachyrhizi* in resistant and susceptible soybean. *Transactions of the British Mycological Society*, *74*, 328–333.
- Kobae, Y., Sekino, T., Yoshioka, H., Nakagawa, T., Martinoia, E., & Maeshima, M. (2006). Loss of AtPDR8, a plasma membrane ABC transporter of *Arabidopsis thaliana*, causes hypersensitive cell death upon pathogen infection. *Plant & Cell Physiology*, *47*, 309–318.
- Kortekamp, A. (2005). Growth, occurrence and development of septa in *Plasmopara viticola* and other members of the *Peronosporaceae* using light- and epifluorescence-microscopy. *Mycological Research*, *109*, 640–648.
- Kortekamp, A. (2006). Expression analysis of defence-related genes in grapevine leaves after inoculation with a host and a non-host pathogen. *Plant Physiology and Biochemistry*, *44*, 58–67.
- Kortekamp, A., & Zyprian, E. (2003). Characterization of *Plasmopara*-resistance in grapevine using in vitro plants. *Journal of Plant Physiology*, *160*, 1393–1400.
- Liu, J. J., & Ekramoddoullah, A. K. M. (2006). The family 10 of plant pathogenesis-related proteins: Their structure, regulation, and function in response to biotic and abiotic

- stresses. *Physiological and Molecular Plant Pathology*, *68*, 3–13.
- Lo, S. C., Hipskind, J. D., & Nicholson, R. L. (1999). cDNA cloning of a sorghum pathogenesis-related protein (PR-10) and differential expression of defense-related genes following inoculation with *Cochliobolus heterostrophus* or *Colletotrichum sublineolum*. *Molecular Plant-Microbe Interactions*, *12*, 479–489.
- Lu, G., Jantasuriyarat, C., Zhou, B., & Wang, G. L. (2004). Isolation and characterization of novel defense response genes involved in compatible and incompatible interactions between rice and *Magnaporthe grisea*. *Theoretical and Applied Genetics*, *108*, 525–534.
- Mauch, F., Mauch-Mani, B., & Boller, T. (1988). Antifungal hydrolases in pea tissue: II. Inhibition of fungal growth by combinations of chitinase and beta-1, 3-glucanase. *Plant Physiology*, *87*, 936–942.
- McGee, J. D., Hamer, J. E., & Hodges, T. K. (2001). Characterization of a PR-10 pathogenesis-related gene family induced in rice during infection with *Magnaporthe grisea*. *Molecular Plant-Microbe Interactions*, *14*, 877–886.
- Mendgen, K., & Hahn, M. (2002). Plant infection and the establishment of fungal biotrophy. *Trends in Plant Science*, *7*, 352–356.
- Monteiro, S., Barakat, M., Picarra-Pereira, M. A., Teixeira, A. R., & Ferreira, R. B. (2003). Osmotin and thaumatin from grape: a putative general defense mechanism against pathogenic fungi. *Phytopathology*, *93*, 1505–1512.
- Moy, P., Qutob, D., Chapman, B. P., Atkinson, I., & Gijzen, M. (2004). Patterns of gene expression upon infection of soybean plants by *Phytophthora sojae*. *Molecular Plant-Microbe Interactions*, *17*, 1051–1062.
- Musetti, R., Marabottini, R., Badiani, M., Martini, M., di Toppi, L. S., Borselli, S., et al. (2007). On the role of H₂O₂ in the recovery of grapevine (*Vitis vinifera* cv. Prosecco) from Flavescence doree disease. *Functional Plant Biology*, *34*, 750–758.
- Okushima, Y., Koizumi, N., Kusano, T., & Sano, H. (2000). Secreted proteins of tobacco cultured BY2 cells: identification of a new member of pathogenesis-related proteins. *Plant Molecular Biology*, *42*, 479–488.
- Panstruga, R. (2003). Establishing compatibility between plants and obligate biotrophic pathogens. *Current Opinion in Plant Biology*, *6*, 320–326.
- Pezet, R., Gindro, K., Viret, O., & Richter, H. (2004). Effects of resveratrol, viniferins and pterostilbene on *Plasmopara viticola* zoospore mobility and disease development. *Vitis*, *43*, 145–148.
- Pina, A., & Errea, P. (2008). Differential induction of phenylalanine ammonia-lyase gene expression in response to in vitro callus unions of *Prunus* spp. *Journal of Plant Physiology*, *165*, 705–714.
- Polesani, M., Bortesi, L., Ferrarini, A., Zamboni, A., Fasoli, M., Zadra, C., et al. (2010). General and species-specific transcriptional responses to downy mildew infection in a susceptible (*Vitis vinifera*) and a resistant (*V. riparia*) grapevine species. *BMC Genomics*, *11*, 117.
- Polesani, M., Desario, F., Ferrarini, A., Zamboni, A., Pezzotti, M., Kortekamp, A., et al. (2008). cDNA-AFLP analysis of plant and pathogen genes expressed in grapevine infected with *Plasmopara viticola*. *BMC Genomics*, *9*, 142.
- Ponchet, M., Panabieres, F., Milat, M. L., Mikes, V., Montillet, J. L., Suty, L., et al. (1999). Are elicitors cryptograms in plant-Oomycete communications? *Cellular and Molecular Life Sciences*, *56*, 1020–1047.
- Restrepo, S., Myers, K. L., del Pozo, O., Martin, G. B., Hart, A. L., Buell, C. R., et al. (2005). Gene profiling of a compatible interaction between *Phytophthora infestans* and *Solanum tuberosum* suggests a role for carbonic anhydrase. *Molecular Plant-Microbe Interactions*, *18*, 913–922.
- Ruepp, A., Zollner, A., Maier, D., Albermann, K., Hani, J., Mokrejs, M., et al. (2004). The FunCat, a functional annotation scheme for systematic classification of proteins from whole genomes. *Nucleic Acids Research*, *32*, 5539–5545.
- Scheel, D. (1998). Resistance response physiology and signal transduction. *Current Opinion in Plant Biology*, *1*, 305–310.
- Sewalt, V. J. H., Ni, W. T., Blount, J. W., Jung, H. G., Masoud, S. A., Howles, P. A., et al. (1997). Reduced lignin content and altered lignin composition in transgenic tobacco down-regulated in expression of L-phenylalanine ammonia-lyase or cinnamate 4-hydroxylase. *Plant Physiology*, *115*, 41–50.
- Smalle, J., & Vierstra, R. D. (2004). The ubiquitin 26 S proteasome proteolytic pathway. *Annual Review of Plant Biology*, *55*, 555–590.
- Stukkens, Y., Bultreys, A., Grec, S., Trombik, T., Vanham, D., & Boutry, M. (2005). NpPDR1, a pleiotropic drug resistance-type ATP-binding cassette transporter from *Nicotiana plumbaginifolia*, plays a major role in plant pathogen defense. *Plant Physiology*, *139*, 341–352.
- Trouvelot, S., Varnier, A. L., Allegre, M., Mercier, L., Baillieux, F., Arnould, C., et al. (2008). A beta-1, 3 glucan sulfate induces resistance in grapevine against *Plasmopara viticola* through priming of defense responses, including HR-like cell death. *Molecular Plant-Microbe Interactions*, *21*, 232–243.
- Unger, S., Buche, C., Boso, S., & Kassemeyer, H. H. (2007). The Course of Colonization of Two Different *Vitis* Genotypes by *Plasmopara viticola* Indicates Compatible and Incompatible Host-Pathogen Interactions. *Phytopathology*, *97*, 780–786.
- Van den Berg, N., Berger, D. K., Hein, I., Birch, P., Wingfield, M. J., & Viljoen, A. (2007). Tolerance in banana to Fusarium wilt is associated with early up-regulation of cell wall-strengthening genes in the roots. *Molecular Plant Pathology*, *8*, 333–341.
- Van den Berg, N., Crampton, B. G., Hein, I., Birch, P., & Berger, D. K. (2004). High-throughput screening of suppression subtractive hybridization cDNA libraries using DNA microarray analysis. *Biotechniques*, *37*, 818–824.
- van Loon, L. C., Rep, M., & Pieterse, C. M. (2006). Significance of inducible defense-related proteins in infected plants. *Annual Review of Phytopathology*, *44*, 135–162.
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., et al. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology*, *3*, 7.

- Velculescu, V. E., Zhang, L., Vogelstein, B., & Kinzler, K. W. (1995). Serial analysis of gene expression. *Science*, *270*, 484–487.
- Vergne, E., Ballini, E., Marques, S., Sidi Mammar, B., Droc, G., Gaillard, S., et al. (2007). Early and specific gene expression triggered by rice resistance gene Pi33 in response to infection by ACE1 avirulent blast fungus. *The New Phytologist*, *174*, 159–171.
- Werner, S., Steiner, U., Becher, R., Kortekamp, A., Zyprian, E., & Deising, H. B. (2002). Chitin synthesis during in planta growth and asexual propagation of the cellulosic oomycete and obligate biotrophic grapevine pathogen *Plasmopara viticola*. *FEMS Microbiology Letters*, *208*, 169–173.
- Wielgoss, A., & Kortekamp, A. (2006). Comparison of PR1 expression in grapevine cultures after inoculation with a host- and a non-host pathogen. *Vitis*, *45*, 9–13.
- Wong, H. L., Sakamoto, T., Kawasaki, T., Umemura, K., & Shimamoto, K. (2004). Down-regulation of metallothionein, a reactive oxygen scavenger, by the small GTPase OsRac1 in rice. *Plant Physiology*, *135*, 1447–1456.
- Yazaki, K. (2006). ABC transporters involved in the transport of plant secondary metabolites. *FEBS Letters*, *580*, 1183–1191.