

The genome of the Gram-positive metal- and sulfate-reducing bacterium *Desulfotomaculum reducens* strain MI-1

Pilar Junier,^{1†} Thomas Junier,² Sheila Podell,³
David R. Sims,⁴ John C. Detter,⁴
Athanasios Lykidis,⁵ Cliff S. Han,⁴
Nicholas S. Wigginton,^{1‡} Terry Gaasterland³ and
Rizlan Bernier-Latmani^{1*}

¹Environmental Microbiology Laboratory, Ecole Polytechnique Fédérale de Lausanne, Lausanne 1015, Switzerland.

²Computational Evolutionary Genomics Group, University of Geneva, Geneva 1211, Switzerland.

³Marine Biological Research Division, Scripps Institution of Oceanography, La Jolla, CA 92037, USA.

⁴DOE Joint Genome Institute, Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA.

⁵DOE-Joint Genome Institute, Genome Biology Program, Walnut Creek, CA 94598, USA.

Summary

Spore-forming, Gram-positive sulfate-reducing bacteria (SRB) represent a group of SRB that dominates the deep subsurface as well as niches in which resistance to oxygen and desiccation is an advantage. *Desulfotomaculum reducens* strain MI-1 is one of the few cultured representatives of that group with a complete genome sequence available. The metabolic versatility of this organism is reflected in the presence of genes encoding for the oxidation of various electron donors, including three- and four-carbon fatty acids and alcohols. Synteny in genes involved in sulfate reduction across all four sequenced Gram-positive SRB suggests a distinct sulfate-reduction mechanism for this group of bacteria. Based on the genomic information obtained for sulfate reduction in *D. reducens*, the transfer of electrons to the sulfite and APS reductases is proposed to take place via the quinone pool and heterodisulfide reductases respectively. In addition,

both H₂-evolving and H₂-consuming cytoplasmic hydrogenases were identified in the genome, pointing to potential cytoplasmic H₂ cycling in the bacterium. The mechanism of metal reduction remains unknown.

Introduction

The genus *Desulfotomaculum* contains sulfate-reducing bacteria (SRB) that form heat-, oxygen- and desiccation-resistant endospores (Widdel and Bak, 1992). Because of their ability to form spores, members of this genus thrive in habitats where anoxic conditions are not maintained permanently and where desiccation occasionally occurs. As a result, *Desulfotomaculum* and the closely related genus *Desulfosporosinus* tend to be the dominant SRB in subsurface environments disturbed by human activities, such as mines, as well as sites where the groundwater level is subject to significant seasonal fluctuations. Phylogenetic studies of the bacterial communities in deep mines in Japan (Ishii *et al.*, 2000) and South Africa (Moser *et al.*, 2003) illustrate the importance of *Desulfotomaculum* spp. in those environments. A separate investigation 4 km deep in a South African mine revealed that species of *Desulfotomaculum* and *Methanobacterium* dominate the microbial communities in those environments (Moser *et al.*, 2005). Recently, the relevance of *Desulfotomaculum* and other Gram-positive spore-forming bacteria, particularly *Desulfosporosinus* and *Clostridium* has come to the fore for the bioremediation of U(VI)-contaminated sites and it appears as though these organisms could play an important role at these sites following electron donor amendments (Chang *et al.*, 2001; Madden *et al.*, 2007; N'Guessan *et al.*, 2008).

Desulfotomaculum reducens MI-1 was isolated from marine sediments heavily contaminated with chromium from Mare Island Naval Shipyard (Tebo and Obratzsova, 1998). It can use a wide range of organic compounds as electron donors, including short-chain fatty acids such as propionate, butyrate and valerate, alcohols such as methanol, ethanol, n-propanol and n-butanol as well as lactate, pyruvate and glucose. More importantly, it can use a large number of electron acceptors for growth: sulfate,

*For correspondence. E-mail rizlan.bernier-latmani@epfl.ch; Tel. (+41) 21 693 5001; Fax (+41) 21 693 6205. Present addresses: †Laboratory of Microbiology, University of Neuchâtel, Neuchâtel 2000, Switzerland; ‡Science, 1200 New York Avenue NW, Washington DC, USA.

thiosulfate, dithionite, as well as elemental sulfur. It is one of the few SRB able to grow by coupling the oxidation of organic compounds to the reduction of Fe(III) to Fe(II) (Tebo and Obraztsova, 1998).

As is typical of many SRB, *D. reducens* ferments pyruvate. However, a recent characterization of *D. reducens* has unveiled unusual metabolic properties. The most striking result is that spores of *D. reducens* are able to catalyse the reduction of U(VI) when H₂, a product of pyruvate fermentation, is provided as an electron donor and when cell-free medium collected after cell growth (spent medium) is added (Junier *et al.*, 2009). Under these conditions, the presence of competing electron acceptors such as nitrate or sulfate does not affect U(VI) reduction (Junier *et al.*, 2010). This is significant due to the extensive nitrate contamination found concomitantly with U(VI) in various US Department of Energy (DOE) sites and the prevalence of nitrate outcompeting U(VI) as an electron acceptor for many microorganisms (Elias *et al.*, 2003; Luo *et al.*, 2005).

Desulfotomaculum reducens MI-1 is an excellent example of a microorganism to be investigated in more detail because its unusual metabolic characteristics make it effectively both a sulfate- and a metal-reducing bacterium. In addition, it is one of the few Gram-positive SRB for which a complete genome sequence is available. The other two are *Desulfotomaculum acetoxidans* (a close relative to *D. reducens*) and Candidatus *Desulforudis audaxviator* that was sequenced from DNA obtained from South African mine (Chivian *et al.*, 2008). It is one of only three SRB with a genome sequence available and able to couple growth to Fe(III) reduction. The other two are *Desulfotalea psychrophila* (Knoblauch *et al.*, 1999) and *Desulfobulbus propionicus* (Holmes *et al.*, 2004).

Eighteen draft and finished SRB genome sequences are available to date and were used for comparison: seven *Desulfovibrio* spp. (*Dv. desulfuricans* G20, *Dv. desulfuricans* ATCC 27774, *Dv. piger*, *Dv. salexigens*, *Dv. vulgaris* Miyazaki, *Dv. vulgaris* DP4, *Dv. vulgaris* Hildenborough), *Desulfonatronospira thiodismutans*, *Desulfohalobium retbaense*, *Desulfococcus oleovorans*, *Desulfobacterium autotrophicum*, *Desulfobulbus propionicus*, *Desulfomicrobium baculatum*, *Desulfotalea psychrophila*, Candidatus *Desulforudis audaxviator*, *Desulfonispora thiosulfatigenes* and two *Desulfotomaculum* spp. (*D. reducens* and *D. acetoxidans*). The latter four are Gram-positive.

In this article, we present the complete genome sequence of *D. reducens* strain MI-1. Using manual annotation, metabolic information as well as the comparison to genomes of the eighteen above-mentioned SRB, we describe general features of the *D. reducens* genome including genetic components involved in central metabolic pathways, sulfate reduction, hydrogen metabolism and metal reduction.

Results and discussion

General genome features

The finished genome of *D. reducens* consists of a single circular chromosome, 3 608 104 base pairs (bp) long, with a G+C content of 43% (Table 1 and Fig. 1). 71 tRNAs were identified. In addition to the 69 standard tRNAs, two are for nonstandard aminoacids: pseudouridine (dred_R0046) and selenocysteine (dred_R0069). The genome contains eight almost identical copies of the three rRNAs (5S, 16S and 23S) but in addition, we found 16S and 5S rRNA genes that were slightly longer than the ones pertaining to operons (1629 bp instead of 1612 bp for 16S rRNA and 205 bp instead of 117 bp for 5S rRNA). A combination of automated annotation and extensive manual curation predicted 3324 protein-coding sequences (CDS), of which, 2334 (70%) were assigned functions.

Central and electron donor metabolism

Central metabolic pathways such as the Embden-Meyerhoff-Parnas (glycolysis) pathway and the reductive pentose phosphate pathway are present. The glycolysis pathway (Fig. S1) and the reductive pentose phosphate pathway (Fig. S2) are complete. A partial reductive citric acid cycle is also present that allows the transformation of oxaloacetate to succinate via malate and fumarate and succinyl-coA to isocitrate via α -ketoglutarate. However, it is unclear whether the succinate to succinyl-coA step is possible because the succinyl-coA synthetase normally

Table 1. *Desulfotomaculum reducens* strain MI-1 genome statistics.

Characteristic	Value
Chromosome size (bp)	3 608 104
G+C ratio (%)	43.28
Coding density (%)	85.5
No. of predicted protein coding gene	3 324
No. of predicted proteins unique to <i>D. reducens</i> (%)	515 (15.5)
Number of rRNA operons	8
Number of tRNA genes	71
No. of predicted proteins with putative function (%)	2 334 (70.2)
No. of predicted proteins with unknown function (%)	990 (29.8)
Genes coding transmembrane proteins	510
BLASTP comparison against completed microbial genome database (No. of top hits)	
<i>Firmicutes</i>	2 553
<i>Clostridia</i>	2 375
<i>Bacilli</i>	178
<i>Deltaproteobacteria</i>	50
<i>Syntrophomonadaceae</i>	32

% GC statistics are for protein coding sequences only. Criteria for BLASTP match inclusion: (i) BLASTP *e*-value $1e - 5$ or lower; (ii) alignment covering > 70% of both query and subject.




Fig. 1. Circular representation of the genome of *Desulfotomaculum reducens* strain MI-1. From the outside inwards: Genes on forward strand (blue) and genes on reverse strand (black) and tRNA genes (green), genes with best blast hit from *D. acetoxidans* (red), genes with best blast hit from *Desulfitobacterium hafniense* Y51 (blue), hypothetical proteins (brown), transposases and integrases (green), and G+C content (purple).

active in this pathway (E.C. 6.2.1.5) is absent from the *D. reducens* genome (Fig. S3). Without this enzyme, the partial citric acid pathway does not allow the formation of α -ketoglutarate from oxaloacetate, suggesting that glutamate metabolism is lacking in *D. reducens*. In fact, the organism does not grow in the absence of yeast extract, suggesting that it is a fastidious bacterium (Fig. S4).

A characteristic of *D. reducens* is its ability to utilize a large number of electron donors: lactate, pyruvate, glucose, short-chain fatty acids such as propionate, butyrate and valerate, and alcohols such as methanol, ethanol, n-propanol or n-butanol (Tebo and Obratzsova, 1998). Fig. 2 shows the pathways involved in the oxidation of butyrate (green) and ethanol (purple).

Pyruvate is an intermediate in the oxidation of several electron donors (propionate, lactate, glucose) in addition to being itself an electron donor and fermentative substrate. The pathways for the metabolism of propionate, lactate and glucose via pyruvate are shown in yellow in Fig. 2. An interesting feature of the *D. reducens* genome is that it encodes genes for all three potential pyruvate transformation mechanisms (Fig. S5; Fig. 2) (White, 2000). Decarboxylation by the pyruvate dehydrogenase (PDH) (dred_1893) yields acetate, CO_2 and NADH;

decarboxylation by the pyruvate-ferredoxin oxidoreductase (PFOR) (dred_0047-50) yields acetate, CO_2 and H_2 and decarboxylation by the pyruvate-formate lyase (PFL) (dred_2750-53) yields acetate and formate. Experimental evidence for activity of the former two is available (Junier *et al.*, 2009; Junier *et al.*, 2010). In order to produce H_2 from the fermentation of pyruvate, an H_2 -evolving hydrogenase is involved (see hydrogen metabolism section).

Acetate is the product of the oxidation of all electron donors because *D. reducens* is an incomplete oxidizer. Thus, acetyl-CoA is also an intermediate in many pathways. Surprisingly, we were unable to identify a phosphotransacetylase to catalyse the formation of acetyl-phosphate from acetyl-CoA in the genome of *D. reducens* (Fig. S5) despite comparing the phosphotransacetylase from *D. acetoxidans* to the entire *D. reducens* genome using BLASTP. Nonetheless, an acetate kinase (E.C. 2.7.2.1; dred_2094) was identified and is predicted to catalyse the transformation of acetyl-phosphate to acetate with the production of ATP. We also identified an acetyl-CoA ligase (E.C. 6.2.1.13; dred_2081) that produces acetate and ATP from acetyl-CoA directly and that is likely to be involved in energy production from the acetyl-CoA intermediate (Fig. S5).




Fig. 2. Metabolic reconstruction of *D. reducens* based on known growth substrates and metabolic capacities. Arrows indicate metabolic flows: dashed lines represent putative electron flow. The sole protein shown in dark blue contains selenocysteine. Abbreviations are: DH, dehydrogenase; H₂ase, hydrogenase; DsrMK, menaquinol oxidizing complex; DsrABC, sulfite reductase; NrfHA, nitrite reductase; Hdr, heterodisulfide reductase; MvhD, methyl viologen reducing hydrogenase, delta subunit; Sat, methyl sulfate adenyl transferase; QmoAB, Hdr- and Mvh-containing protein complex; FeS, iron sulfur protein; Fd, ferredoxin, EtfAB, electron-transfer flavoprotein; OR, oxidoreductase; PPI, pyrophosphate.

Two complete non-identical copies of the aerobic-type carbon monoxide dehydrogenase (CoxMSL) are present (dred_2774-6 and dred_1502-4). This protein is a dimer of heterotrimers: it contains a molybdoprotein, a flavoprotein and an Fe-S protein and catalyses the oxidation of CO coupled to the reduction of an electron acceptor (Schubel *et al.*, 1995). This aerobic-type carbon monoxide dehydrogenase (CODH) was recently identified in the complete genomes of two other anaerobes: *Carboxydothemus hydrogeniformans* (Luo *et al.*, 2005) and *Moorella thermoacetica* (moth_1958-60) (Pierce *et al.*, 2008). This suggests that the aerobic-type CODH may function with other electron acceptors than O₂ (e.g., sulfate). We tested *D. reducens* for its ability to utilize CO as an electron donor in the presence of sulfate as an electron acceptor with or without an additional electron donor (lactate). We found that CO was inhibitory to sulfate reduction in the presence of lactate and did not support sulfate reduction in the

presence of CO alone (Fig. S6) suggesting no direct involvement of CODH in bulk CO oxidation. Two studies (Voordouw, 2002; Pereira *et al.*, 2008) suggest that CO may be produced during sulfate reduction in *Desulfovibrio vulgaris*. The transcriptomic study (Pereira *et al.*, 2008) suggests that CO may be produced during pyruvate decarboxylation and is consistent with CO serving as a physiological electron donor for energy-producing processes at low concentrations. Thus, even if the aerobic type CODH does not support growth with CO as an electron donor and CO is toxic to cells at the concentrations considered, CODH may be involved in the cycling of low levels of CO within the cell during sulfate reduction (see sulfate section).

Sulfate catabolism

Unlike other types of respiration where the terminal reductases are located on the cytoplasmic (or the outer)

membrane, sulfate reduction is biochemically challenging because of the cytoplasmic localization of the terminal reductases [adenosine phosphosulfate (APS) reductase and sulfite reductase]. This means that the direct translocation of protons across the membrane coupled to electron transport cannot take place. A model for the pathway of dissimilatory sulfate reduction involving 'hydrogen cycling' was developed for *Desulfovibrio vulgaris* (Odom and Peck, 1981). The general scheme is predicated on the production of H₂ that diffuses freely into the periplasm and carries with it both electrons and protons. In the H₂ cycling model, hydrogen gas in the periplasm donates electrons to periplasmic hydrogenases that transfer them to periplasmic c₃ cytochromes, releasing protons into the periplasm and generating a proton motive force. The hydrogen cycling model has been expanded to include formate and CO as potential intermediates shuttling electrons from the cytoplasm to the periplasm and is termed 'redox cycling' (Pereira *et al.*, 2008). Similarly to hydrogen cycling, periplasmic formate dehydrogenases oxidize formate and release protons, electrons and CO₂ into the periplasm (Pereira *et al.*, 2008). Cytoplasmic CO is converted to periplasmic CO₂ and H₂ via a membrane-spanning CODH-hydrogenase complex (Voordouw, 2002). Subsequently, the electrons that have been accumulated in the periplasmic c₃ cytochrome pool are transferred back into the cytoplasm to the APS and sulfite reductases through transmembrane complexes (e.g., Qmo, Rnf, Dsr, Hmc) (Heidelberg *et al.*, 2004).

Desulfotomaculum reducens is a Gram-positive SRB and, thus, lacks a periplasmic space as well as periplasmic proteins. As a result, the electron flow during sulfate reduction must differ from that in *Dv. vulgaris*. In this section, we propose a mechanism by which *D. reducens* obtains energy from sulfate reduction in the absence of a periplasm (shown in red in Fig. 2).

Activation of sulfate to APS. The sulfate adenylyltransferase (dred_0635) releases pyrophosphate during the conversion of sulfate to APS. A fourteen transmembrane helix (TMH) proton-translocating pyrophosphatase (Fig. 2) is encoded in the genome of *D. reducens* (dred_2985) as well as in the genome of all three other Gram-positive SRB. In contrast, only a soluble pyrophosphatase is found in Gram-negative SRB. The transmembrane pyrophosphatase may allow the translocation of protons during the cleavage of the pyrophosphate diphosphate bond, suggesting that the activation of sulfate may contribute to the establishment of a proton motive force. Its presence in all four Gram-positive SRB sequenced to date is consistent with a strategy to obtain energy from sulfate reduction in the absence of periplasmic proteins.

Reduction of APS to sulfite. Adjacent to the sulfate adenylyltransferase is the APS reductase (dred_0636-7) that is homologous to APS reductases from other SRB. However, the alpha subunit (dred_0637) of the APS reductase in *D. reducens* was predicted to be membrane-anchored and outside-facing using the transmembrane helix (TMH) prediction algorithm TMHMM (<http://www.cbs.dtu.dk/services/TMHMM>). Out of the eighteen SRB genomes, *D. reducens* is the only one for which the APS reductase alpha subunit is predicted to be membrane-anchored (with a probability > 85%). Note that the *Desulfonispota* sp. genome sequence does not cover this region and thus, the APS reductase from that organism was not included in this analysis. Bioinformatic evidence of the membrane-anchored localization for the *D. reducens* APS reductase will need to be confirmed by biochemical analyses as single transmembrane helix predictions are not as reliable as multi-helix predictions. However, there is a precedent for this observation. Immunocytochemical evidence shows the association of the APS reductase with the cytoplasmic membrane in the thermophilic SRB *Desulfovibrio thermophilus* whereas it confirms the cytoplasmic localization of the protein in *Dv. vulgaris* and *Dv. gigas* (Kremer *et al.*, 1988).

Associated with the adenylyltransferase and the APS reductase are a heterodisulfide reductase (Hdr) (dred_0638) and the fusion protein of a heterodisulfide reductase and the delta subunit of the methyl-viologen-reducing [Ni-Fe] hydrogenase (Mvh) (dred_0639). The latter two ORFs (dred_0638 and dred_0639) are conserved in all SRB sequenced to date (Fig. 3) and represent two of the three genes of the QmoABC complex in *Dv. vulgaris*. QmoABC is a transmembrane complex that, in *Dv. vulgaris*, allows the transfer of electron from the periplasmic c₃ cytochrome pool to the terminal reductases in the cytoplasm (Haveman *et al.*, 2004). While the gamma subunit (QmoC) of QmoABC includes TMHs in all Gram-negative SRB, the three examples of Gram-positive SRB (*D. reducens*, *D. acetoxidans* and Candidatus *Desulfurudis* – the *Desulfonispota* sp. draft sequence does not include the appropriate region) lack TMHs. In *D. reducens*, QmoC is completely absent. Because the Qmo operon is adjacent to the APS reductase, it has been suggested that it may be involved in transferring electrons from periplasmic c-type cytochromes to this cytoplasmic reductase in *Dv. desulfuricans* (Pires *et al.*, 2003; Haveman *et al.*, 2004). The fact that there is no transmembrane subunit in the Qmo complex in the three Gram-positive SRB for which sequence is available suggests that the Qmo complex is unlikely to catalyse transmembrane electron transfer in those bacteria. Nonetheless, it is possible that QmoAB in *D. reducens* serves as an electron donor to APS reductase if electrons are donated to the complex from a cytoplasmic source such




Fig. 3. *aprBA* and *qmoABC* organization for all available SRB genome sequences. *aprBA*, beta and alpha subunit of the APS reductase; *qmoABC*, alpha, beta and gamma subunits of the quinone-interacting membrane-bound oxidoreductase. QmoC is the transmembrane subunit but is absent in *D. reducens* and soluble in *D. acetoxidans* and *D. audaxviator*. *cspL*, cold shock protein; *hyp*, hypothetical protein; *hdrB*, heterodisulfide reductase; *qmoAB*, genes encoding a heterodisulfide reductase and a methyl-viologen reducing hydrogenase protein complex.

as electron transfer flavoproteins (EtfAB) or heterodisulfide reductase (Hdr) (Fig. 2).

Reduction of sulfite to sulfide. The sulfite reductase (dred_3186-7) in *D. reducens* is homologous to sulfite reductases from all other SRB. In *Desulfovibrio* spp. and presumably other Gram-negative SRB, a transmembrane complex, the Dsr complex, is postulated to transfer periplasmic electrons to the sulfite reductase (Heidelberg *et al.*, 2004). The equivalent complex is dubbed Hme complex in *Desulfobacterium autotrophicum* (Strittmatter *et al.*, 2009). The Dsr complex is a five-protein complex (DsrMKJOP) (Pires *et al.*, 2006) whose encoding genes show almost perfect synteny across all 14 sequenced Gram-negative SRB (Fig. 4). In *Dv. desulfuricans*, DsrM is an integral membrane cytochrome *b* and DsrK is a soluble protein homologous to the HdrD subunit of the heterodis-

ulfide reductase (Pires *et al.*, 2006). Using the Ortholog Neighborhood Viewer tool in JGI-IMG (<http://img.jgi.doe.gov>) for *Dv. vulgaris* Hildenborough's *dsrMKJOP* against all SRB genomes, we were able to identify orthologous genes for *dsrMK* but not for *dsrJOP* in *D. reducens* (Fig. 4). These genes were annotated as the gamma subunit of a nitrate reductase (dred_3199) and a putative reductase (dred_3198). The gamma subunit of the nitrate reductase (dred_3199, *dsrM*) is a *b*-type cytochrome and is predicted to have five transmembrane helices and to pertain to a protein family (PF02665) that receives electrons from the quinone pool. While there is some similarity between dred_3198-9 and *dsrMK* from *Dv. vulgaris* Hildenborough, it is clear that there is a dichotomy between these genes and associated proteins in Gram-positive bacteria and their counterparts in Gram-negative bacteria. In fact, a BLAST analysis of the




Fig. 4. Gene organization for the *dsr* operon for all available genome sequences. *cbiA*, cobyrinic acid synthase; *cutA*, copper toxicity protein; *FeS*, iron sulfur protein; *nfi*, endonuclease V; *asp*, aspartyl aminopeptidase. Unlabelled genes are hypothetical proteins.

membrane-spanning protein encoded by *dred_3199* yields no hits for Gram-negative SRB. It has been suggested that the Dsr complex is made up of two modules DsrMK and DsrJOP and that the DsrMK module might be involved in menaquinol oxidation as well as the reduction of a cytoplasmic substrate (Pires *et al.*, 2006). That description of the activity of DsrMK, in the absence of DsrJOP, is consistent with its potential electron transport role (from the quinone pool to a cytoplasmic protein) in *D. reducens* and other Gram-positive SRB (Fig. 2).

Adjacent to *dsrMK*, arranged as part of what appears to be the same operon, the *D. reducens* genome (and that of

the other three sequenced Gram-positive SRB) encodes a soluble protein (*dred_3197*) that corresponds to the gamma subunit of the dissimilatory sulfite reductase (DsrC) (Fig. 4). In *D. vulgaris*, DsrC has recently been implicated in serving as an electron shuttle between the membrane-bound DsrMKJOP complex and the cytoplasmic DsrAB sulfite reductase (Oliveira *et al.*, 2008). In that study, DsrC was found to interact with DsrK (the cytoplasmic catalytic subunit) and to form a persulfide-containing intermediate that once dissociated from DsrAB was released as sulfide. We hypothesize that, in *D. reducens* and other Gram-positive SRB, DsrC may similarly transfer




Fig. 5. Seven *hdr* loci in *D. reducens*. *etfAB*, electron transfer flavoprotein; *paaH*, 3-hydroxybutyryl-CoA dehydrogenase; *crt*, 3-hydroxybutyryl-CoA dehydratase; *bcd*, butyryl-CoA dehydrogenase; *atoB*, acetyl-CoA C-acetyltransferase; *cat2*, 4-hydroxybutyrate coenzyme A transferase; *bioB*, biotin synthetase; *bacA*, undecaprenyl-diphosphatase; FeS, iron-sulfur protein; *mvhD*, methyl viologen hydrogenase, delta subunit; MFS, major facilitator superfamily protein; OR, oxidoreductase; *tmh*, transmembrane helices-containing protein; *hdrA*, *hdrB*, *hdrC*, *hdrD*, *hdrF*, different heterodisulfide reductase subunits; *hdrF1*, transmembrane heterodisulfide reductase; *hdrA1*, NADH-dependent heterodisulfide reductase. Locus I is dred_0137 to dred_0148; Locus II is dred_0633 to dred_0639 (includes *qmoAB* which are dred_0638-9); locus III is dred_0358 to dred_0354; locus IV is dred_1325 to dred_1330; locus V is dred_1777 to dred_1786; locus VI is dred_0427 to dred_0433 and locus VII is dred_0689 to dred_0690. The asterisk "*" indicates a selenocysteine-containing protein.

electrons from DsrK to DsrAB, which in turn, is involved in the reduction of sulfite to sulfide. The overall electron transfer would then take place from the quinone pool to DsrM, DrsK, DsrC and finally DsrAB (Fig. 2).

Electron flow to APS and sulfite reductases. Other transmembrane electron transfer complexes present in *Dv. vulgaris* (e.g., HmcABCDEF or RnfABCDGE) are not found in *D. reducens* or other Gram-positive SRB, which is consistent with the presence of a distinct sulfate reduction mechanism in Gram-positive SRB. In *D. reducens*, the electrons are generated via the oxidation of organic substrates and transit via NADH, H₂, formate, CO as well as various reduced soluble electron transport proteins such as ferredoxin (dred_2206) and electron transfer flavoproteins (EtfAB) (dred_1778-9, dred_1538-9, dred_0572-3 and dred_0367-8) (Fig. 2).

Formate is oxidized via formate dehydrogenase (dred_1112-19) that includes a ten-TMH transmembrane subunit (dred_1116) annotated as a polysulfide reductase and that is part of a protein family (PF03916) that is known to participate in electron exchange with the quinone pool. The oxidation of formate to CO₂ could be coupled to sodium translocation (Fig. 2) as the last ORF in the operon (dred_1119) encodes a sodium-dependent transporter. This ORF is slightly upregulated during lactate-

dependent sulfate reduction (data not shown), suggesting the formation of formate as an intermediate of lactate oxidation. The electrons are presumably passed on to the quinone pool. Similarly, NADH is oxidized by a quinone-interacting, proton-translocating NADH dehydrogenase (DH) (dred_2036-46) that also passes electron to the quinone pool (Fig. 2). From the quinone pool, the electrons are transferred to DsrM and on to sulfite reductase, as suggested above.

Heterodisulfide reductase. The genome of *D. reducens* encodes numerous heterodisulfide reductases (Hdr) found in seven loci (Fig. 5). A recent analysis of the genome of *Desulfobacterium autotrophicum* HRM2 showed nine *hdr* loci in that organism. Many of the cytoplasmic Hdr in *Db. autotrophicum* were presumed to be involved in (i) the transfer of electron from transmembrane complexes to the sulfite reductase, (ii) the reduction of the disulfide bond of the DsrABC complex, and (iii) the reduction of ferredoxin (Strittmatter *et al.*, 2009). In *D. reducens*, the *hdr* loci include several gene configurations: *hdrA* associated with *mvhD* (loci I, II and IV), *hdrA*, *hdrB* and *hdrC* in close proximity (loci II, III, IV), a membrane-spanning *hdrF* associated with *etfAB* and with *hdrD* (locus V) and *hdrD* alone (loci VI and VII) (Fig. 5). Locus I consists of two identical repeats of the same

sequence (dred_0137 to dred_0141 and dred_0143 to dred_0148) separated by a gene encoding a protein of unknown function (dred_0142).

In two loci (II and IV), the presence of all three subunits of Hdr (HdrABC) along with the delta subunit of an Mvh (a [NiFe] H₂ase) is reminiscent of the Mvh : Hdr complex in the methanogen *Methanothermobacter thermoautotrophicus* (Setzke *et al.*, 1994). This complex is thought to transfer electrons from H₂ to an electron acceptor interacting with the Hdr (Mander *et al.*, 2004). The main difference between the methanogen and *D. reducens* is the absence of the other subunits of Mvh : *Methanothermobacter marburgensis* has three (Stojanowic *et al.*, 2003), including two (MvhA and MvhG) that correspond to hydrogenase modules conserved in all [NiFe] H₂ases, but *D. reducens* only has the delta subunit (MvhD) that is unique to Mvh (Table S1). The delta subunit has been implicated in the transfer of electrons from Mvh to Hdr (Stojanowic *et al.*, 2003). By analogy, electrons derived from a soluble cytoplasmic protein could be transferred to MvhD and on to HdrA (locus I) or HdrABC (locus IV) and to the APS reductase (Fig. 2).

Locus V combines the 4-TMH membrane-spanning *hdrF* (dred_1777) with electron-transfer flavoproteins (dred_1778-9) and an *hdrD* subunit (dred_1783) (Fig. 5). It is reminiscent of HdrDE in *Methanosarcina barkeri* (Heiden *et al.*, 1994) but lacks HdrE, the membrane-anchoring *b*-type cytochrome. In *Methanosarcina mazei*, this complex is a proton pump that is part of the electron transport chain and receives electrons from hydroxyphenazine and uses them to reduce the CoenzymeM–CoenzymeB complex (Ide *et al.*, 1999; Baumer *et al.*, 2000). In *D. reducens*, the membrane-spanning domain pertains to HdrF and it is likely that this subunit receives electrons from the quinone pool and passes them on to the electron-transfer flavoproteins (Fig. 2). It is also possible that proton–translocation occurs during the electron transfer (Fig. 2).

If the current proposed scheme for sulfate reduction in *D. reducens* (Fig. 2) is confirmed, it would represent a significant departure from the accepted model of ‘redox cycling’ for sulfate reduction by Gram-negative SRB. In the proposed scheme (shown in red in Fig. 2), electrons from electron donors (such as lactate or butyrate) would be transferred to the Qmo complex from intermediate electron-carrying proteins such as heterodisulfide reductase and/or electron transfer flavoproteins, and Qmo would pass the electrons on to APS reductase, which in turn, would reduce APS to sulfite. Electrons could also be transferred from electron donors to electron-carrying compounds such as H₂, formate and NADH. Those electrons would be transferred to the quinone pool by proteins such as the membrane-bound hydrogenase, formate DH and NADH DH. From the quinone pool, elec-

trons could be transferred to the DsrC subunit of the sulfite reductase via the DsrMK complex and sulfite reduced to sulfide. The generation of a proton motive force, which relies on the presence of periplasmic H₂ in Gram-negative bacteria, would occur via the translocation of protons through pyrophosphatase, formate DH, membrane-bound H₂ase, the quinone pool as well as NADH DH. Given that many of the proteins proposed to be involved in sulfate reduction in *D. reducens* have homologues in the other sequenced Gram-positive SRB (i.e., pyrophosphatase, DsrMK, QmoAB, HdrABC, MvhD), *D. reducens* may serve as a model for sulfate reduction in Gram-positive SRB.

Hydrogen metabolism

In general, the consumption and the evolution of H₂ are mediated by hydrogenases (H₂ases), redox metalloenzymes that catalyse the reversible reaction $H_2 \leftrightarrow 2H^+ + 2e^-$ (Vignais *et al.*, 2001; Vignais and Colbeau, 2004; Meyer, 2007). These enzymes have been classified as either [FeFe], [NiFe] or [NiFeSe] H₂ases according to the composition of their metal sites.

The *D. reducens* genome encodes six hydrogenases (Table 2). This number of hydrogenases is comparable with that in the genomes of organisms that, in contrast to *D. reducens*, are able to conserve energy by using H₂ as a source of electrons. For instance, the genome of *Dv. vulgaris* in which sulfate reduction depends on H₂ oxidation encodes six hydrogenases (Heidelberg *et al.*, 2004), and that of *Geobacter sulfurreducens*, which is able to couple H₂ oxidation to growth with Fe(III) as an electron acceptor, has four (Methe *et al.*, 2003). All six H₂ases in *D. reducens* are Fe-containing hydrogenases and four of them are encoded by multi-locus operons and represent trimeric H₂ases (Table 2). The other two (dred_1440 and dred_1794) are single-gene hydrogenases.

Trimeric Fe-containing H₂ases. Three (dred_3290-92, dred_1651-53 and dred_1654-56) of the four multimeric [FeFe] H₂ases are cytoplasmic, trimeric hydrogenases composed of a catalytic subunit and two additional subunits and are encoded by non-identical copies of the operon. The genes encoding the catalytic subunit (dred_3290, dred_1651 and dred_1654) contain several FeS binding sites in addition to the active site. The additional subunits are homologous to two peripheral subunits of NADH-ubiquinone oxidoreductase, indicating that these are NAD(P)(H) dependent H₂ases. These trimeric hydrogenases resemble the H₂-evolving [FeFe] H₂ases of the hyperthermophilic bacterium *Thermotoga maritima* that is able to grow by fermenting carbohydrates (Verhagen *et al.*, 1999). The three H₂ases were found to be upregulated during sulfate reduction with lactate as an

Table 2. List of hydrogenases identified in the *D. reducens* genome, their characteristics and their expression level during pyruvate fermentation or lactate-dependent sulfate reduction.

Locus dred_	Product name	Type	Organization	Localization	Size (aa)	Expression level	
						Fermentation	Respiration
1440	Hydrogenase	FeFe	Monomeric	Unknown	429	0.63 ± 0.19	1.47 ± 0.61
1794	Hydrogenase	FeFe	Monomeric	Unknown	462	3.37 ± 0.24	0.35 ± 0.03
0461	Hydrogenase	FeFe	Trimeric	Membrane-bound	381	0.94 ± 0.20	1.02 ± 0.04
0462					261	1.14 ± 0.17	0.95 ± 0.06
0463 ^a					520	1.39 ± 0.12	0.97 ± 0.03
1651 ^a	Hydrogenase	FeFe	Trimeric	Cytoplasmic	593	0.16 ± 0.09	1.88 ± 0.10
1652					569	0.12 ± 0.08	2.06 ± 0.09
1653					158	0.12 ± 0.06	2.34 ± 0.07
1654 ^a	Hydrogenase	FeFe	Trimeric	Cytoplasmic	659	0.48 ± 0.06	19.75 ± 0.79
1655					627	0.47 ± 0.16	12.76 ± 0.74
1656					163	0.42 ± 0.10	11.09 ± 0.47
3290 ^a	Hydrogenase	FeFe	Trimeric	Cytoplasmic	594	0.39 ± 0.02	2.16 ± 0.54
3291					575	0.09 ± 0.02	3.06 ± 1.05
3292					177	0.11 ± 0.02	2.97 ± 1.08

a. Indicates the catalytic domain-containing subunit in trimeric hydrogenases.

electron donor but not during pyruvate fermentation (Table 2), suggesting that H₂ production may be mediated by these trimeric [FeFe] H₂ases and coupled to H₂ consumption (no H₂ is detected in the medium) during lactate oxidation.

The fourth trimeric [FeFe] hydrogenase (dred_0461-3) is membrane-bound and is encoded by an operon that includes a FeS-cluster-containing component (dred_0462) and a putative hydrogenase cytochrome *b* subunit (dred_0461) that is predicted to contain ten transmembrane helices. This H₂ase is homologous to a potentially H₂-consuming hydrogenase in *Desulfitobacterium hafniense* Y51 (Nonaka *et al.*, 2006). Thus, this H₂ase could be involved in the transfer of electrons from H₂ to the quinone pool and – by virtue of its membrane localization – transfer protons to the outside (Fig. 2).

Monomeric Fe-containing H₂ases. These two H₂ases (dred_1440 and dred_1794) are upregulated (Table 2) either during pyruvate fermentation (dred_1794), suggesting the evolution of H₂, or during sulfate reduction (dred_1440) suggesting the consumption of H₂.

Phylogenetic study of the catalytic subunit. A phylogenetic study of the catalytic subunit of the six [FeFe] hydrogenases (Fig. 6) reveals that there are two major clusters. One corresponds to fast-evolving hydrogenases (the branches are long and there is more evolutionary distance between adjacent species) and the other to slow-evolving hydrogenases (shorter branches with less evolutionary distance between adjacent species). It is readily apparent that the trimeric H₂ases are in the slow-evolving branch whereas their monomeric counterparts are in the fast-evolving branch. This is consistent with the need for evolution in concert of the three subunits for the trimeric

case whereas the monomeric H₂ases can evolve independently.

The overall picture that emerges from the above sections is that the trimeric cytoplasmic [FeFe] H₂ases are responsible for the production of H₂ during sulfate reduction, the trimeric [FeFe] membrane-spanning H₂ase consumes H₂ and translocates H⁺ and the monomeric [FeFe] H₂ases either produce (dred_1794) or consume (dred_1440) H₂.

Metal reduction

Vegetative cells of *D. reducens* are able to reduce soluble Fe(III) with pyruvate, lactate or butyrate but not H₂ as an electron donor, in a process that supports growth (Fig. S7). This is an unusual ability in SRB as there are only four other SRB known to couple Fe(III) reduction to growth: *D. psychrophila* (Knoblauch *et al.*, 1999), *Desulfosporosinus lacus* (Ramamoorthy *et al.*, 2006), *Desulfosporomusa polytropa* (Sass *et al.*, 2004) and *Desulfobulbus propionicus* (Holmes *et al.*, 2004). Furthermore, vegetative cell suspensions (Tebo and Obratzsova, 1998) as well as spores (Junier *et al.*, 2009) of *D. reducens* were shown to reduce U(VI) with butyrate and H₂ as an electron donor, respectively.

The biochemical mechanism of microbially mediated iron reduction has been studied in some detail in two bacterial genera: *Geobacter* and *Shewanella*. In *Geobacter* spp., direct contact between Fe(III) oxides and bacterial cells is required and the production of conductive ‘nanowires’ facilitates the transfer of electrons to the solid phase (Reguera *et al.*, 2005). The transfer of electrons from the NADH dehydrogenase to the terminal electron acceptor during iron catabolism requires five *c*-type cytochromes and a type IV pilus (Weber *et al.*, 2006).




Fig. 6. Neighbour-joining phylogenetic tree of Fe-only hydrogenases based on alignments of the H cluster domain. Clades consisting entirely of the same genus were condensed to a single leaf. The branches corresponding to the different Fe-only hydrogenase genes in *Desulfotomaculum reducens* are boxed in blue for trimeric hydrogenases and in red for monomeric hydrogenases.

Shewanella oneidensis MR-1 also produces conductive appendages (Gorby *et al.*, 2006) and also requires *c*-type cytochromes for soluble and insoluble Fe(III) reduction (Beliaev *et al.*, 2001). In addition, the use of endogenous and exogenous soluble extracellular electron shuttles has been well documented to mediate the reduction of

Fe(III)-oxides in both genera (Newman and Kolter, 2000; Nevin and Lovley, 2002).

As for microbial U(VI) reduction, in *S. oneidensis*, the pathways for iron and U(VI) reduction are similar and involve several *c*-type cytochromes (Beliaev *et al.*, 2001; Bencheikh-Latmani *et al.*, 2005). U(VI) reduction in *G.*

sulfurreducens involves other *c*-type cytochromes than those involved in Fe(III) reduction (Shelobolina *et al.*, 2007).

In contrast, the genome of *D. reducens* is strikingly cytochrome-poor. The only *c*-type cytochrome present is a triheme cytochrome *c*₅₅₂ that is encoded by two genes (*dred_0700* and *dred_0701*). This is an integral membrane *c*-type cytochrome annotated as the two-subunit nitrite reductase (NrfHA). This protein is likely to transfer electrons from the quinone pool to an unidentified cytoplasmic or extracellular electron acceptor. Evidence for such a process rests in the small subunit (NrfH) that is predicted (via BLAST analysis) to interact with menaquinol. Even though NrfHA is annotated as a nitrite reductase, it is unlikely to be involved in nitrite reduction. Analysis of mRNA from a *D. reducens* culture amended with nitrite during sulfate reduction shows that this gene has a higher qualitative level of expression in the absence than in the presence of nitrite (Fig. 7C). This is in sharp contrast to the effect of nitrite on the expression of the physiological nitrite reductase in *Dv. vulgaris* for which genes encoding nitrite reductase were highly upregulated (Haveman *et al.*, 2004). Thus, the low nitrite reduction activity observed could be due to the sulfite reductase that is known to have some nitrite reductase activity in *Dv. vulgaris* (Haveman *et al.*, 2004). We hypothesize that, in *D. reducens*, NrfHA may play a role in electron transfer directly, or indirectly via the quinone pool, to electron acceptors other than nitrite, for example U(VI) (shown in blue, Fig. 2) or soluble Fe(III).

Inspection of electron micrographs of *D. reducens* after U(VI) reduction suggests that the process is associated with the cell wall because the accumulation of U(IV) occurs both inside and outside the cell membrane (Fig. 8). NrfHA is an outside-facing, transmembrane *c*-type cytochrome. Due to its localization, we hypothesize that this protein may be involved in U(VI) reduction.

Several studies have considered U(VI) and Fe(III) reduction by the two SRB *Dv. vulgaris* and *Dv. desulfuricans*. The mechanism of reduction was found to involve both periplasmic (Payne *et al.*, 2002) and cytoplasmic (Li and Krumholz, 2009) proteins. Uranium reduction in *Dv. desulfuricans* with H₂ as an electron donor was found to involve the transfer of electrons from a periplasmic hydrogenase via the periplasmic *c*₃ cytochromes to U(VI) (Payne *et al.*, 2002). In *Dv. vulgaris* Hildenborough, Fe(III) and U(VI) reduction was found to also involve periplasmic H₂ases and *c*₃-type cytochromes (Elias *et al.*, 2004). A more recent study has shown that it is a periplasmic [FeFe] H₂ase, not a [NiFe] H₂ase, that is involved in the reduction of soluble Fe(III) (Park *et al.*, 2008). By analogy, the H₂-consuming, membrane-bound [FeFe] hydrogenase in *D. reducens* (Table 2) could be involved in U(VI) reduction. Paralleling the mechanism described in *Des-*




Fig. 7. Effect of nitrite on growth of and sulfate reduction by *D. reducens*.

A. Cell growth as OD₆₀₀ (closed symbols and solid lines) sulfate (closed symbols and dashed lines) and nitrite (open symbols and dashed lines) reduction. The addition of 0.1 mM nitrite inhibits growth and sulfate reduction, but the cells recover after nitrite is removed from solution. In contrast, 0.2 mM nitrite shuts down growth and sulfate reduction.

B. Nitrite reductase operon.

C. Reverse transcription PCR results for *dsrAB* and *nrfHA* after addition of nitrite. NrfA is expressed in the absence of nitrite but to a lesser extent in its presence. NrfH is not detectable in any of the cases considered.

ulfovibrio spp., the hydrogenase could be transferring electrons from H₂ to the *c*-type cytochrome NrfHA, which would serve as the terminal reductase (shown in blue, Fig. 2).

In addition to the H₂ases, a cytoplasmic thioredoxin, a thioredoxin reductase and associated oxidoreductase were found to participate in U(VI) reduction in the cytoplasm of *Dv. desulfuricans* G20 (Li and Krumholz, 2009). In an effort to determine whether a similar mechanism could be involved in *D. reducens*, we evaluated the pres-




Fig. 8. Electron micrograph of thin sections of a *D. reducens* culture grown fermentatively with pyruvate in the presence of 100 μ M U(VI) and stained with Sato lead. Scale bar is 1 μ m.

ence of thioredoxins in its genome. There are three ORFs annotated as thioredoxins in the *D. reducens* genome (dred_0762, dred_0904 and dred_2669), but only one (dred_2669) is associated with a thioredoxin reductase (dred_2670). A third gene (dred_2668) that includes nine TMH and has no known function is present as part of this apparent operon. This thioredoxin could possibly be involved in U(VI) reduction, but no direct evidence for this is available at the present time.

Conclusions

Desulfotomaculum reducens is one of two cultivated Gram-positive SRB for which a complete genome is available. Due to the absence of a periplasm, the mechanism of sulfate-reduction in this SRB does not correspond to that identified in *Desulfovibrio vulgaris* and likely involves the transfer of electrons directly from the quinone pool to the DsrMK complex and to the sulfite reductase (DsrABC). In addition, it is proposed that the cytoplasmic QmoAB complex transfers electrons from cytoplasmic proteins to the APS reductase. Other proteins such as heterodisulfide reductases and electron-transfer flavoproteins are likely involved in electron transfer and we propose that they may provide electrons to the Qmo complex. However, a detailed mechanism cannot be surmised by genome analysis. Finally, a number of proton-

translocating transmembrane proteins – notably a pyrophosphatase conserved in all four Gram-positive SRB sequenced to date – suggest the production of a proton motive force without periplasmic oxidation of H_2 . Because many of the proteins involved in the *D. reducens*-specific sulfate-reduction pathway have homologues in the genomes of other Gram-positive SRB, *D. reducens* may represent a good model for sulfate-reduction in Gram-positive SRB.

Experimental procedures

DNA extraction and purification

Genomic DNA was extracted from an overnight culture of *D. reducens* that was grown in Widdel medium (Widdel and Bak, 1992) with sulfate (10 mM) as an electron acceptor and lactate (20 mM) as an electron donor. DNA extraction was carried out using CTAB (hexadecyltrimethyl ammonium bromide). The cells were treated with 100 mg ml^{-1} lysozyme, followed by proteinase K (10 mg ml^{-1}) and the product amended with a solution of 0.3 M CTAB and 0.7 M NaCl and a chloroform:isoamyl alcohol (24:1) solution. After centrifugation of the mixture, the supernatant was transferred to a new tube and amended with a phenol:chloroform:isoamyl alcohol (25:24:1) solution. After a second centrifugation, the supernatant was transferred and the DNA precipitated with isopropanol. A final treatment with RNase A (Qiagen, Valencia, CA) was used to remove residual RNA.

Sequencing

The genome of *D. reducens* MI-1 was sequenced at the Joint Genome Institute (JGI) using a combination of 3 kb, 8 kb and 40 kb (fosmid) DNA libraries. All general aspects of library construction and sequencing performed at the JGI can be found at <http://www.jgi.doe.gov/>. Draft assemblies were based on 40 920 total reads. All three libraries provided 11 \times coverage of the genome. The Phred/Phrap/Consed software package (<http://www.phrap.com>) was used for sequence assembly and quality assessment (Ewing and Green, 1998; Ewing *et al.*, 1998; Gordon *et al.*, 1998). After the shotgun stage, reads were assembled with parallel phrap (High Performance Software, LLC). Possible mis-assemblies were corrected with Dupfinisher (Han and Chain, 2006) or transposon bombing of bridging clones (Epicentre Biotechnologies, Madison, WI, USA). Gaps between contigs were closed by editing in Consed, custom primer walk or PCR amplification (Roche Applied Science, Indianapolis, IN, USA). A total of 2 634 additional reactions were necessary to close gaps and to raise the quality of the finished sequence. The completed genome sequences of *D. reducens* contains 43 940 reads, achieving an average of 12-fold sequence coverage per base with an error rate less than 1 in 100 000.

Annotation

Genes were identified using a combination of Critica (Badger and Olsen, 1999) and Glimmer (Delcher *et al.*, 1999), fol-

lowed by a round of manual curation, which resulted in adjustments of the start codons and insertion of missed genes and pseudogenes. tRNAs were predicted using the tRNAscan-SE tool (Lowe and Eddy, 1997). Automatic product name assignment was made based on the results obtained from searches against curated databases. Signal peptides were identified using the SignalP 3.0 (Bendtsen *et al.*, 2004) and TMHMM (Krogh *et al.*, 2001) at default values. Manual curation of the automatic annotation occurred within the IMG-ER (<http://merced.jgi-psf.org/cgi-bin/er/main.cgi>) system. The sequence data described here have been deposited in GenBank (CP000612).

Phylogenetic analysis of Fe-Fe hydrogenases

The hydrogenase phylogeny was based on the H cluster, which is shared by all members of the hydrogenase family (Vignais *et al.*, 2001). The position of the H cluster in one sequence was determined by scanning for Pfam PF02906 (H cluster) with Prosite. The sequences were then aligned with Muscle (Edgar, 2004), and the alignment was trimmed so as to keep only the H cluster. A phylogenetic tree was then constructed using BioNJ (Gascuel, 1997), using a eukaryote (*Culex quinquefasciatus*) as an outgroup. Clades consisting entirely of the same genus were condensed to a single leaf.

Nitrite reduction as a detoxification strategy

To test nitrite reduction, a culture was grown in WLP medium supplemented with yeast extract, 0.05%; NaHCO₃, 30 mM; 1,4-piperazinediethane sulfonic acid disodium salt monohydrate (PIPES), 20 mM; sodium sulfate, 10 mM, lactic acid 20 mM. The culture was incubated for 9.5 h until the onset of growth and sulfate reduction. Aliquots of the culture (20 ml) were transferred to 50 ml serum bottles previously flushed with N₂. The cultures were then supplemented with 0, 0.1 or 0.2 mM of sodium nitrite. Growth was quantified by measuring optical density at 600 nm. Samples for SO₄²⁻ and NO₂⁻ were collected, filtered (PVDF 0.2 µm filters) and analysed by ion chromatography (DX-500, Dionex, Sunnyvale, CA, USA) using an IonPac AS12A column and a bicarbonate (30 mM) eluent. A sample for RNA extraction was collected at 3.5 h of incubation after the addition of nitrite. RNA was extracted using the RNAeasy Qiagen Kit with on-column DNase treatment. The quantity and quality of the RNA were evaluated with a Nanodrop spectrophotometer (Thermo Scientific). Reverse transcription PCR (RT-PCR) was carried out for the following genes: 16S rRNA, *nrfH* and *nrfA*. For RT-PCR primers specific for *D. reducens* were designed using Primer-BLAST: 16S rRNA, DR140 (5'-TAG ACC GGG ATA ACA GCT G-3') and DR842 (5'-ATA CCC GCA ACA CCT AGC AC-3'); *nrfH*, *nrfHF* (5'-CAT TAT GGA TCC CTG GGT TG -3') and *nrfHR* (5'-GTC CTG ACC ACG GTC ATT CT-3'), and *nrfA*, *nrfAF* (5'-AGC CCC GGA GTC ACT TTT AT-3') and *nrfAR* (5'-CAT GAC ACT GGG CAC ATA CC-3'). First-strand synthesis was carried out using SuperScript III from Invitrogen. PCR was carried out using NEB *Taq* DNA polymerase.

Hydrogenase expression data

For expression analysis Widdel medium was used. The medium was dispensed (100 ml) into 200 ml glass serum

bottles and autoclaved. The following solutions were added from sterile anaerobic stocks (final concentration): yeast extract, 0.05%; NaHCO₃, 30 mM; 1,4-piperazinediethane sulfonic acid disodium salt monohydrate (PIPES), 20 mM. For fermentative growth only 20 mM of pyruvic acid was added to the medium. For sulfate respiration 20 mM sulfate (as sodium sulfate) and 20 mM lactic acid were included in the medium. The final pH of the medium was 7.2 ± 0.2. Growth in the cultures was monitored by OD₆₀₀. The experiments were run in four biological replicates. The cultures were sampled (16 ml) for RNA extraction at mid-exponential phase. Samples were collected in an RNase-free 50 ml Falcon tube by centrifugation at 7000 *g* for 7 min and resuspended in 400 µl of 3 mg ml⁻¹ lysozyme in TE buffer (pH 8.0) and mixed by vortexing. After digestion for 10 min at room temperature, 1.4 ml of Buffer RLT (Qiagen) containing freshly added 0.01% v/v β-mercaptoethanol was added to the sample and mixed vigorously by vortexing. The homogenized cell lysates were stored at -80°C. After all the samples were collected, the cell lysates were thawed for 15 min at 37°C in a water bath to dissolve salts. The samples were separated into four equal aliquots of 450 µl. RNA extraction was carried out as described above. RNA from the four aliquots was combined and precipitated using 0.1 volume of 1 M sodium acetate and 2.5 volume cold 95% ethanol and incubated over-night at -20°C. RNA was collected by centrifugation for 15 min at 12 000 *g* at 4°C and washed with 75% ethanol. RNA was dried at 37°C for 20 min and resuspended in 30 µl RNase-free water. RNA concentration and quality were re-measured using the Nanodrop and the Bioanalyzer (Agilent). A total of 20 µg of RNA per sample was sent to Nimblegen Roche for cDNA synthesis and hybridization onto custom-designed 4x77K microarrays. Normalized RMA signals were used for the analysis using the software GeneSpring GX v7.3. Per gene normalization was applied when the data were loaded into a customized one-colour experiment in the analysis software. Expression data were extracted using the option 'advanced find gene'. Average values were calculated from the four biological replicates.

Electron microscopy

Samples for electron microscopy were prepared by fixing the cells in glutaraldehyde and dehydrating them sequentially in pure grade Ethanol (Fluka). The fixed dehydrated cells were then pelleted and immobilized in LR-white resin (EMS, Hatfield, PA, USA) that was polymerized at 60°C. This procedure was carried out inside an anaerobic chamber to prevent sample oxidation. Thin sections of the resin were cut using a microtome, placed on copper grids (Quantifoil Micro Tools GmbH, Jena) and stained with Sato lead. The samples were observed in a FEI CM20 microscope (Eindhoven, Netherlands). Images were recorded on a Gatan 797 slow scan CCD camera (1024 pixels × 1024 pixels × 14 bits) and processed with the Gatan Digital Micrograph 3.11.0 software (Gatan, Pleasanton, CA, USA).

Access to the genome is provided by the DOE Joint Genome Institute (JGI) at the Integrated Microbial Genome (IMG) site: <http://img.jgi.doe.gov/cgi-bin/pub/main.cgi>

Acknowledgements

We acknowledge funding from the Swiss National Science Foundation through Grant No. 33100A0-112337. In addition, this publication was made possible by grant number ES010337 from the National Institute of Environmental Health Sciences (NIEHS), NIH. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIEHS, NIH. The work conducted by the US Department of Energy Joint Genome Institute is supported by the Office of Science of the US Department of Energy under contract No. DE-AC02-05CH11231. T.G.'s contributions to computational aspects of this work were supported in part by the US National Science Foundation grant number 0626678. Finally, we thank three anonymous reviewers whose constructive comments lead to a marked improvement of this manuscript.

References

- Badger, J.H., and Olsen, G.J. (1999) CRITICA: coding region identification tool invoking comparative analysis. *Mol Biol Evol* **16**: 512–524.
- Baumer, S., Ide, T., Jacobi, C., Johann, A., Gottschalk, G., and Deppenmeier, U. (2000) The F420-H₂ dehydrogenase from *Methanosarcina mazei* is a redox-driven proton pump closely related to NADH dehydrogenases. *J Biol Chem* **275**: 17968–17973.
- Beliaev, A.S., Saffarini, D.A., McLaughlin, J.L., and Hunnicutt, D. (2001) MtrC, an outer membrane decahaem c cytochrome required for metal reduction in *Shewanella putrefaciens* MR-1. *Mol Microbiol* **39**: 722–730.
- Bencheikh-Latmani, R., Williams, S.M., Haucke, L., Criddle, C.S., Wu, L., Zhou, J., and Tebo, B.M. (2005) Global Transcriptional Profiling of *Shewanella oneidensis* MR-1 During Cr(VI) and U(VI) Reduction. *Appl Environ Microbiol* **71**: 7453–7460.
- Bendtsen, J.D., Nielsen, H., von Heijne, G., and Brunak, S. (2004) Improved prediction of signal peptides: signalP 3.0. *J Mol Biol* **340**: 783–795.
- Chang, Y., Peacock, A.D., Long, P., Stephen, J.R., McKinley, J.P., Macnaughton, S.J., et al. (2001) Diversity and characterization of sulfate-reducing bacteria in groundwater at a uranium mill tailings site. *Appl Environ Microbiol* **67**: 3149–3160.
- Chivian, D., Brodie, E.L., Alm, E.J., Culley, D.E., Dehal, P.S., DeSantis, T.Z., et al. (2008) Environmental genomics reveals a single-species ecosystem deep within earth. *Science* **322**: 275–278.
- Delcher, A.L., Harmon, D., Kasif, S., White, O., and Salzberg, S.L. (1999) Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* **27**: 4636–4641.
- Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797.
- Elias, D.A., Krumholz, L.R., Wong, D., Long, P.E., and Suflita, J.M. (2003) Characterization of microbial activities and U reduction in a shallow aquifer contaminated by uranium mill tailings. *Microb Ecol* **46**: 83–91.
- Elias, D.A., Suflita, J.M., McInerney, M.J., and Krumholz, L.R. (2004) Periplasmic cytochrome c₃ of *Desulfovibrio vulgaris* is directly involved in H₂-mediated metal but not sulfate reduction. *Appl Environ Microbiol* **70**: 413–420.
- Ewing, B., and Green, P. (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res* **8**: 186–194.
- Ewing, B., Hillier, L., Wendl, M.C., and Green, P. (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res* **8**: 175–185.
- Gascuel, O. (1997) BIONJ: an improved version of the NJ algorithm based on a simple model of sequence data. *Mol Biol Evol* **14**: 685–695.
- Gorby, Y.A., Yanina, S., McLean, J.S., Rosso, K.M., Moyles, D., Dohnalkova, A., et al. (2006) Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms. *Proc Natl Acad Sci USA* **103**: 11358–11363.
- Gordon, D., Abajian, C., and Green, P. (1998) Consed: a graphical tool for sequence finishing. *Genome Res* **8**: 195–202.
- Han, C.S., and Chain, P. (2006) Finishing repeat regions automatically with Dupfinisher. In *Proceedings of 2006 International Conference on Bioinformatics & Computational Biology*. Arabnia, H.R., and Valafar, H. (eds). Las Vegas, NV, USA: CSREA Press, pp. 141–146.
- Haveman, S.A., Greene, E.A., Stilwell, C.P., Voordouw, J.K., and Voordouw, G. (2004) Physiological and gene expression analysis of inhibition of *Desulfovibrio vulgaris* Hildenborough by nitrite. *J Bacteriol* **186**: 7944–7950.
- Heidelberg, J.F., Seshadri, R., Haveman, S.A., Hemme, C.L., Paulsen, I.T., Kolonay, J.F., et al. (2004) The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Nat Biotechnol* **22**: 554–559.
- Heiden, S., Hedderich, R., Setzke, E., and Thauer, R.K. (1994) Purification of a two-subunit cytochrome-b-containing heterodisulfide reductase from methanol-grown *Methanosarcina barkeri*. *Eur J Biochem* **221**: 855–861.
- Holmes, D.E., Bond, D.R., and Lovley, D.R. (2004) Electron transfer by *Desulfobulbus propionicus* to Fe(III) and graphite electrodes. *Appl Environ Microbiol* **70**: 1234–1237.
- Ide, T., Baumer, S., and Deppenmeier, U. (1999) Energy-conservation by the H₂:heterodisulfide oxidoreductase from *Methanosarcina mazei* Go1: identification of two proton-translocating segments. *J Bacteriol* **181**: 4076–4080.
- Ishii, K., Takii, S., Fukunaga, S., and Aoki, K. (2000) Characterization by denaturing gradient gel electrophoresis of bacterial communities in deep groundwater at the Kamaishi Mine, Japan. *J Gen Appl Microbiol* **46**: 85–93.
- Junier, P., Frutschi, M., Wigginton, N.S., Schofield, E.J., Bargar, J.R., and Bernier-Latmani, R. (2009) Metal reduction by spores of *Desulfotomaculum reducens*. *Environ Microbiol* **11**: 3007–3017.
- Junier, P., Suvorova, E.I., and Bernier-Latmani, R. (2010) Effect of competing electron acceptors on the reduction of U(VI) by *Desulfotomaculum reducens* *Geomicrobiol J* (in press).
- Knoblauch, C., Sahm, K., and Jorgensen, B.B. (1999) Psychrophilic sulfate-reducing bacteria isolated from permanently cold Arctic marine sediments: description of *Desulfofrigus oceanense* gen. nov., sp. nov., *Desulfofrigus*

- fragile* sp nov., *Desulfofaba gelida* gen. nov., sp nov., *Desulfotalea psychrophila* gen. nov., sp nov and *Desulfotalea arctica* sp nov. *Int J Syst Bacteriol* **49**: 1631–1643.
- Kremer, D.R., Veenhuis, M., Fauque, G., Peck, H.D., Legall, J., Lampreia, J., et al. (1988) Immunocytochemical localization of APS reductase and bisulfite reductase in three *Desulfovibrio* species. *Arch Microbiol* **150**: 296–301.
- Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E.L.L. (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**: 567–580.
- Li, X., and Krumholz, L.R. (2009) Thioredoxin is involved in U(VI) and Cr(VI) reduction in *Desulfovibrio desulfuricans* G20. *J Bacteriol* **191**: 4924–4933.
- Lowe, T.M., and Eddy, S.R. (1997) tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* **25**: 955–964.
- Luo, J., Cirpka, O.A., Wu, W.M., Fienen, M.N., Jardine, P.M., Mehlhorn, T.L., et al. (2005) Mass-transfer limitations for nitrate removal in a uranium-contaminated aquifer. *Environ Sci Technol* **39**: 8453–8459.
- Madden, A.S., Smith, A.C., Balkwill, D.L., Fagan, L.A., and Phelps, T.J. (2007) Microbial uranium immobilization independent of nitrate reduction. *Environ Microbiol* **9**: 2321–2330.
- Mander, G.J., Pierik, A.J., Huber, H., and Hedderich, R. (2004) Two distinct heterodisulfide reductase-like enzymes in the sulfate-reducing archaeo *Archaeoglobus profundus*. *Eur J Biochem* **271**: 1106–1116.
- Methe, B.A., Nelson, K.E., Eisen, J.A., Paulsen, I.T., Nelson, W., Heidelberg, J.F., et al. (2003) Genome of *Geobacter sulfurreducens*: metal reduction in subsurface environments. *Science* **302**: 1967–1969.
- Meyer, J. (2007) [FeFe] hydrogenases and their evolution: a genomic perspective. *Cell Mol Life Sci* **64**: 1063–1084.
- Moser, D.P., Onstott, T.C., Fredrickson, J.K., Brockman, F.J., Balkwill, D.L., Drake, G.R., et al. (2003) Temporal shifts in the geochemistry and microbial community structure of an ultradeep mine borehole following isolation. *Geomicrobiol J* **20**: 517–548.
- Moser, D.P., Gihring, T.M., Brockman, F.J., Fredrickson, J.K., Balkwill, D.L., Dollhopf, M.E., et al. (2005) *Desulfotomaculum* and *Methanobacterium* spp. dominate a 4-to 5-kilometer-deep fault. *Appl Environ Microbiol* **71**: 8773–8783.
- N'Guessan, A.L., Vrionis, H.A., Resch, C.T., Long, P.E., and Lovley, D.R. (2008) Sustained removal of uranium from contaminated groundwater following stimulation of dissimilatory metal reduction. *Environ Sci Technol* **42**: 2999–3004.
- Nevin, K.P., and Lovley, D.R. (2002) Mechanisms for Fe(III) oxide reduction in sedimentary environments. *Geomicrobiol J* **19**: 141–159.
- Newman, D.K., and Kolter, R. (2000) A role for excreted quinones in extracellular electron transfer. *Nature* **405**: 94–97.
- Nonaka, H., Keresztes, G., Shinoda, Y., Ikenaga, Y., Abe, M., Naito, K., et al. (2006) Complete genome sequence of the dehalorespiring bacterium *Desulfitobacterium hafniense* Y51 and comparison with *Dehalococcoides ethenogenes* 195. *J Bacteriol* **188**: 2262–2274.
- Odom, J.M., and Peck, H.D. (1981) Hydrogen cycling as a general mechanism for energy coupling in the sulfate-reducing bacteria *Desulfovibrio* sp. *FEMS Microbiol Lett* **12**: 47–50.
- Oliveira, T.F., Vorrhein, C., Matias, P.M., Venceslau, S.S., Pereira, I.A.C., and Archer, M. (2008) The crystal structure of desulfovibrio vulgaris dissimilatory sulfite reductase bound to dsrC provides novel insights into the mechanism of sulfate respiration. *J Biol Chem* **283**: 34141–34149.
- Park, H.S., Lin, S., and Voordouw, G. (2008) Ferric iron reduction by *Desulfovibrio vulgaris* Hildenborough wild type and energy metabolism mutants. *Anton Leeuw Int J G* **93**: 79–85.
- Payne, R.B., Gentry, D.M., Rapp-Giles, B.J., Casalot, L., and Wall, J.D. (2002) Uranium reduction by *Desulfovibrio desulfuricans* Strain G20 and a cytochrome c3 mutant. *Appl Environ Microbiol* **68**: 3129–3132.
- Pereira, P.M., He, Q., Valente, F.M.A., Xavier, A.V., Zhou, J.Z., Pereira, I.A.C., and Louro, R.O. (2008) Energy metabolism in *Desulfovibrio vulgaris* Hildenborough: insights from transcriptome analysis. *Anton Leeuw Int J G* **93**: 347–362.
- Pierce, E., Xie, G., Barabote, R.D., Saunders, E., Han, C.S., Detter, J.C., et al. (2008) The complete genome sequence of *Moorella thermoacetica* (f. Clostridium thermoacetikum). *Environ Microbiol* **10**: 2550–2573.
- Pires, R.H., Lourenco, A.I., Morais, F., Teixeira, M., Xavier, A.V., Saraiva, L.M., and Pereira, I.A.C. (2003) A novel membrane-bound respiratory complex from *Desulfovibrio desulfuricans* ATCC 27774. *BBA-Bioenergetics* **1605**: 67–82.
- Pires, R.H., Venceslau, S.S., Morais, F., Teixeira, M., Xavier, A.V., and Pereira, I.A.C. (2006) Characterization of the *Desulfovibrio desulfuricans* ATCC 27774 DsrMKJOP complex – A membrane-bound redox complex involved in the sulfate respiratory pathway. *Biochemistry* **45**: 249–262.
- Ramamoorthy, S., Sass, H., Langner, H., Schumann, P., Kroppenstedt, R.M., Spring, S., et al. (2006) *Desulfosporosinus lacus* sp nov., a sulfate-reducing bacterium isolated from pristine freshwater lake sediments. *Int J Syst Evol Microbiol* **56**: 2729–2736.
- Reguera, G., McCarthy, K.D., Mehta, T., Nicoll, J.S., Tuominen, M.T., and Lovley, D.R. (2005) Extracellular electron transfer via microbial nanowires. *Nature* **435**: 1098–1101.
- Sass, H., Overmann, J., Rutters, H., Babenzien, H.D., and Cypionka, H. (2004) *Desulfosporomusa polytropica* gen. nov., sp nov., a novel sulfate-reducing bacterium from sediments of an oligotrophic lake. *Arch Microbiol* **182**: 204–211.
- Schubel, U., Kraut, M., Morsdorf, G., and Meyer, O. (1995) Molecular characterization of the gene cluster *coxmsl* encoding the molybdenum containing carbon monoxide dehydrogenase in *Oligotropha carboxidovorans*. *J Bacteriol* **177**: 2197–2203.
- Setzke, E., Hedderich, R., Heiden, S., and Thauer, R.K. (1994) H₂: heterodisulfide oxidoreductase complex from *Methanobacterium thermoautotrophicum*: composition and properties. *Eur J Biochem* **220**: 139–148.
- Shelobolina, E.S., Coppi, M.V., Korenevsky, A.A., DiDonato, L.N., Sullivan, S.A., Konishi, H., et al. (2007) Importance of c-type cytochromes for U(VI) reduction by *Geobacter sulfurreducens*. *BMC Microbiol* **7**: 16.

- Stojanowic, A., Mander, G.J., Duin, E.C., and Hedderich, R. (2003) Physiological role of the F420-non reducing hydrogenase (Mvh) from *Methanothermobacter marburgensis*. *Arch Microbiol* **180**: 194–203.
- Strittmatter, A.W., Liesegang, H., Rabus, R., Decker, I., Amann, J., Andres, S., *et al.* (2009) Genome sequence of *Desulfobacterium autotrophicum* HRM2, a marine sulfate reducer oxidizing organic carbon completely to carbon dioxide. *Environ Microbiol* **11**: 1038–1055.
- Tebo, B.M., and Obratzsova, A.Y. (1998) Sulfate-reducing bacterium grows with Cr(VI), U(VI), Mn(IV), and Fe(III) as electron acceptors. *FEMS Microbiol Lett* **162**: 193–198.
- Verhagen, M.F., O'Rourke, T., and Adams, M.W. (1999) The hyperthermophilic bacterium, *Thermotoga maritima*, contains an unusually complex iron-hydrogenase: amino acid sequence analyses versus biochemical characterization. *BBA-Bioenergetics* **1412**: 212–229.
- Vignais, P.M., and Colbeau, A. (2004) Molecular biology of microbial hydrogenases. *Curr Issues Mol Biol* **6**: 159–188.
- Vignais, P.M., Billoud, B., and Meyer, J. (2001) Classification and phylogeny of hydrogenases. *FEMS Microbiol Rev* **25**: 455–501.
- Voordouw, G. (2002) Carbon monoxide cycling by *Desulfovibrio vulgaris* Hildenborough. *J Bacteriol* **184**: 5903–5911.
- Weber, K.A., Achenbach, L.A., and Coates, J.D. (2006) Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat Rev Microbiol* **4**: 752–764.
- White, D. (2000) *The Physiology and Biochemistry of Prokaryotes*. New York, NY, USA: Oxford University Press.
- Widdel, F., and Bak, F. (1992) Gram-negative mesophilic sulfate-reducing bacteria. In *The Prokaryotes*. Ballows, A., Truper, H.G., Dworkin, M., Harder, W., and Shleifer, K.-H. (eds). Berlin, Germany: Springer, pp. 3352–3378.

Supporting information

Additional supporting information may be found in the online version of this article.

Fig. S1. Genes involved in glycolysis in *D. reducens*.

Fig. S2 Genes involved in the reductive pentose phosphate pathway in *D. reducens*.

Fig. S3 Genes involved in the partial reductive TCA cycle in *D. reducens*.

Fig. S4 Growth of *D. reducens* in the presence and absence of yeast extract.

Fig. S5 Genes involved in pyruvate metabolism in *D. reducens*.

Fig. S6 Inhibition of sulfate reduction in *D. reducens* by CO.

Fig. S7 Fe(III) reduction by *D. reducens*.

Table S1. Genes with homology to the delta subunit of methyl-viologen hydrogenase.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.