


$[(p\text{-Pr}^i\text{C}_6\text{H}_4\text{Me})_4\text{Ru}_4\text{Mo}_4\text{O}_{16}]$: an amphiphilic organoruthenium oxomolybdenum cluster presenting a unique framework geometry

Georg Süss-Fink,* Laurent Plasseraud, Vincent Ferrand and Helen Stoeckli-Evans


Institut de Chimie, Université de Neuchâtel, CH-2000 Neuchâtel, Switzerland

The amphiphilic title compound, easily accessible from (*p*-cymene)ruthenium dichloride dimer and sodium molybdate in aqueous solution, presents an unprecedented $\text{Ru}_4\text{Mo}_4\text{O}_{12}$ framework comprising a central Mo_4O_4 cube with four folded ORuO flaps resembling the sails of a windmill.

The past two decades have witnessed a steadily growing interest in molecules containing both organometallic groups and oxometallic entities,¹ particularly since they provide molecular models for heterogeneous catalysts derived from organometallic complexes adsorbed at metal oxide surfaces.² Organometallic metaloxo clusters contain soft as well as hard metal centers, and hydrophobic as well as hydrophilic ligands. Since the discovery of the first species of this type, $[(\text{C}_5\text{H}_5)\text{TiPW}_{11}\text{O}_{39}]^{4-}$ in 1978,³ this field has been pioneered mainly by the groups of Klemperer,⁴ Isobe,⁵ and Finke.⁶ The combination of low- and high-valent transition metals and the amphiphilic character predispose these molecules also as homogeneous catalysts for oxidation reactions; the catalytic potential of these compounds has been reviewed recently.⁷



Here, we report the synthesis and structure of $[(p\text{-Pr}^i\text{C}_6\text{H}_4\text{Me})_4\text{Ru}_4\text{Mo}_4\text{O}_{16}]$ **1**, a neutral organoruthenium oxomolybdenum cluster which presents an unprecedented geometry of the $\text{Ru}_4\text{Mo}_4\text{O}_{12}$ framework. Compound **1** is easily accessible from (*p*-cymene)ruthenium dichloride dimer and sodium molybdate in aqueous solution (Scheme 1).[†] Orange crystals of correct elemental composition are obtained by crystallization from dichloromethane–toluene. Compound **1** is amphiphilic, that is to say sparingly soluble in both water and aromatic hydrocarbons, and well soluble in polar organic solvents such as dichloromethane or methanol.



Scheme 1

The FAB mass spectrum of **1** shows the molecular ion at m/z 1581 presenting the expected Ru_4Mo_4 isotope pattern. In the ^1H

NMR spectrum (CD_2Cl_2) of **1**, the four *p*-cymene ligands give rise to only one set of signals at δ 1.39 (d, 6, J 7 Hz), 2.27 (s, 3), 2.97 (spt, 1, J 7 Hz), 5.30 (d, 2, J 6 Hz) and 5.37 (d, 2, J 6 Hz), showing all ruthenium positions to be equivalent. In the IR spectrum (KBr) of **1**, the Mo=O stretching vibrations are observed at 921s and 874m cm^{-1} , while the Mo–O–Ru stretches are assigned to the absorptions at 785s, 739s, 642m and 602s cm^{-1} .

Single-crystal X-ray structure analysis[§] of $1 \cdot 2\text{C}_6\text{H}_5\text{Me}$ (orange, rod-like crystals) revealed the molecule to contain a unique $\text{Ru}_4\text{Mo}_4\text{O}_{12}$ framework which can be described as a central Mo_4O_4 cube with four folded ORuO flaps resembling the sails of a windmill. The distortion $[\text{Mo}(2)\text{--O}(2)\text{--Mo}(1) 106.04(12)^\circ, \text{O}(12)\text{--Mo}(1)\text{--O}(2) 71.37(10)^\circ]$ with respect to





Fig. 1 ORTEP¹¹ representation of the molecular structure of **1**. (a) Complete molecule, hydrogen atoms and two toluene molecules (in the crystal) omitted for clarity. (b) $\text{Ru}_4\text{Mo}_4\text{O}_{12}$ framework with labelling scheme. Selected bond distances and angles ($^\circ$): Mo(1)–O(2) 2.122(3), Mo(1)–O(3) 1.804(3), Mo(1)–O(7) 1.706(3), Mo(1)–O(9) 2.073(3), Mo(1)–O(12) 2.365(3), Mo(1)–O(16) 1.798(3). Ru(1)–O(1) 2.088(3), Ru(1)–O(2) 2.089(3), Ru(1)–O(3) 2.094(3); O(12)–Mo(1)–O(9) 72.20(11), Mo(3)–O(12)–Mo(1) 105.31(11), O(7)–Mo(1)–O(16) 103.93(2). Ru(2)–O(9)–Mo(1) 145.40(2), O(10)–Ru(2)–O(11) 80.53(12).

the idealized cubic representation is due to the geometry of the oxygen atoms of the central cube which is more tetrahedral than octahedral (Fig. 1). Eight of the twelve framework oxygen atoms are doubly bridging, while four are quadruply bridging. Each ruthenium atom is coordinated to an η^6 -*p*-cymene ligand [Ru(1)–C₆ ring centroid 1.627 Å], and each molybdenum atom carries a terminal oxo ligand [Mo(1)–O(7) 1.706(3) Å].

The cluster [(C₅Me₅)₄Rh₄Mo₄O₁₆] **2**, reported by Isobe and coworkers,^{5a} which is isoelectronic to **1**, has a completely different structure: the Rh₄Mo₄O₁₂ framework forms a triple cube which is more closed than the open windmill-like structure of **1**. As a consequence, each molybdenum atom carries two terminal oxo ligands, and four of the eight framework oxygen atoms are triply bridging, while the other four are quadruply bridging.



Another Mo₄O₁₆ unit, found in the solid phase Cs₃Mo₄P₃O₁₆, also contains a central Mo₄O₁₆ cube; however, the framework geometry is completely different: the Mo₄O₁₆ core must be considered as four MoO₆ octahedra each sharing three edges with three MoO₆ units giving an array with tetrahedral symmetry.⁷

Interestingly, with ruthenium the reaction seems to work only for the *p*-cymene derivative. Up to now it has not been possible to isolate the benzene and hexamethylbenzene analogues of **1**.

This work was supported by the Fonds National Suisse de la Recherche Scientifique (grant 20-46592.96). A generous loan of ruthenium trichloride hydrate from the Johnson Matthey Technology Centre is gratefully acknowledged.

Footnotes and References

* E-mail: suess-fink@ich.unine.ch

† *Synthesis of 1*: A suspension of [(*p*-Pr^tC₆H₄Me)₂Ru₂Cl₄] (200 mg, 3.27 × 10⁻⁴ mol) in 20 cm³ of water was added dropwise to an aqueous solution (10 cm³) of Na₂MoO₄·2H₂O (790 mg, 3.3 × 10⁻³ mol). The mixture was

stirred for 4 h at 25 °C. After evaporation of the water, the product was extracted with CH₂Cl₂ (20 cm³) and crystallized from dichloromethane–toluene (1:1) to give, after drying *in vacuo*, analytically pure orange crystals (155 mg, 60%).

§ *Crystal structure data for 1*: C₄₀H₅₆Mo₄O₁₆Ru₄·2C₇H₈ (including two molecules of toluene per cluster), triclinic, space group, *P*1̄, *a* = 13.846(2), *b* = 15.210(2), *c* = 16.1038(14) Å, α = 88.124(11), β = 78.938(10), γ = 65.471(11)°, *U* = 3023.7(6) Å³, *Z* = 2, *D_c* = 1.939 g cm⁻³, *T* = 223 K, μ (Mo–K α) = 1.66 mm⁻¹. The data were measured using a Stoe-Siemens AED2 four-circle diffractometer with graphite-monochromated Mo–K α radiation (λ = 0.710 73 Å). 11234 independent reflexions were measured by ω -2 θ scans in the range 2.0 < θ < 25.5°; 9939 were considered observed [*I* > 2 σ (*I*)]. The structure was solved by Patterson and Fourier techniques using the program SHELXS-86⁸ and refined by full-matrix least squares on *F*², using SHELXL-93.⁹ Hydrogen atoms were included in calculated positions and treated as riding atoms using SHELXL-93 default parameters. An empirical absorption correction was applied using DIFABS¹⁰ (transmission factors min., max. 0.722, 1.174). Refinement converged at *R*₁ = 0.0337 and *wR*₂ = 0.0757 (observed data). CCDC 182/532.

- 1 V. W. Day and W. G. Klemperer, *Science*, 1985, **228**, 533.
- 2 K. Isobe and A. Yagasaki, *Acc. Chem. Res.*, 1993, **26**, 524.
- 3 R. K. C. Ho and W. G. Klemperer, *J. Am. Chem. Soc.*, 1978, **100**, 6772.
- 4 H. K. Chae, W. G. Klemperer and V. W. Day, *Inorg. Chem.*, 1989, **28**, 1424; H. K. Chae, W. G. Klemperer, D. E. Paez Loyo, V. W. Day and T. A. Ebersbacher, *Inorg. Chem.*, 1991, **31**, 3187; H. K. Chae, W. G. Klemperer and T. A. Marquart, *Coord. Chem. Rev.*, 1993, **128**, 209.
- 5 (a) Y. Hayashi, K. Toriumi and K. Isobe, *J. Am. Chem. Soc.*, 1988, **110**, 3666; (b) Y. Do, X.-Z. You, C. Zhang, Y. Ozawa and K. Isobe, *J. Am. Chem. Soc.*, 1991, **113**, 5892; (c) Y. Hayashi, Y. Ozawa and K. Isobe, *Inorg. Chem.*, 1991, **30**, 1025; (d) J. T. Park, T. Nishioka, T. Suzuki and K. Isobe, *Bull. Chem. Soc. Jpn.*, 1994, **67**, 1968.
- 6 Y. Lin, K. Nomiya and R. G. Finke, *Inorg. Chem.*, 1993, **32**, 6040; A. Trovarelli and R. G. Finke, *Inorg. Chem.*, 1993, **32**, 6034; M. Pohl and R. G. Finke, *Organometallics*, 1993, **12**, 1453; B. M. Rapko, M. Pohl and R. G. Finke, *Inorg. Chem.*, 1994, **33**, 3625; M. Pohl, D. K. Lyon, N. Mizuno, K. Nomiya and R. G. Finke, *Inorg. Chem.*, 1995, **34**, 1413; K. Nomiya, Ch. Nozaki, M. Kaneko, R. G. Finke and M. Pohl, *J. Organomet. Chem.*, 1995, **505**, 23; M. Pohl, Y. Lin, T. J. R. Weakley, K. Nomiya, M. Kaneko, H. Weiner and R. G. Finke, *Inorg. Chem.*, 1995, **34**, 767.
- 7 C. L. Hill and C. M. Prosser-McCartha, *Coord. Chem. Rev.*, 1995, **143**, 407.
- 8 G. M. Sheldrick, *Acta Crystallogr., Sect. A*, 1990, **46**, 467.
- 9 G. M. Sheldrick, Program for Crystal Structure Refinement, Universität Göttingen, 1993.
- 10 N. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1990, **39**, 158.
- 11 C. K. Johnson, ORTEP, Oak Ridge National Laboratory, Oak Ridge TN, 1976; modified by L. Zolnai and H. Pritzkow, Universität Heidelberg, 1994.

Received in Basel, Switzerland, 2nd May 1997; 7/03035E