

Aqueous oxidation of alcohols catalyzed by artificial metalloenzymes based on the biotin–avidin technology

Christophe M. Thomas, Christophe Letondor, Nicolas Humbert, Thomas R. Ward *

Institut de Chimie, Université de Neuchâtel, Rue Emile-Argand 11, Case Postale 2, CH-2007 Neuchâtel, Switzerland

Received 28 September 2004; received in revised form 29 December 2004; accepted 6 January 2005

Available online 24 March 2005

Abstract

Based on the incorporation of biotinylated organometallic catalyst precursors within (strept)avidin, we have developed artificial metalloenzymes for the oxidation of secondary alcohols using *tert*-butylhydroperoxide as oxidizing agent. In the presence of avidin as host protein, the biotinylated aminosulfonamide ruthenium piano stool complex **1** (0.4 mol%) catalyzes the oxidation of *sec*-phenethyl alcohol at room temperature within 90 h in over 90% yield. Gel electrophoretic analysis of the reaction mixture suggests that the host protein is not oxidatively degraded during catalysis.

Keywords: Artificial metalloenzymes; Ruthenium; Oxidation; Piano stool complex; Biotin–avidin technology

Homogeneous and enzymatic catalysis are in many respects complementary [1–4]. By incorporating an organometallic complex into a host protein, we aim at creating artificial metalloenzymes with properties reminiscent both of enzymes and of homogeneous catalysts. With this goal in mind, we have recently described the incorporation of achiral biotinylated rhodium–diphosphine complexes into (strept)avidin ((strept)avidin refers to either avidin or streptavidin, used as a host protein) [5]. These artificial metalloenzymes were used as catalysts for the enantioselective hydrogenation of *N*-protected dehydroaminoacids (ee up to 96%). The perspective of improving the performance of such artificial metalloenzymes using combined chemical- and genetic optimization strategies (i.e., chemogenetic) provides the opportunity to focus on more challenging reactions. For example, it may be interesting to test the power of such methodology for the oxidation of alcohols. Along similar lines, several groups have recently reported the

creation of artificial metalloenzymes for the enantioselective oxidation of sulfides to sulfoxides [6–8].

The oxidation of alcohols to the corresponding aldehydes and ketones is one of the most important functional group transformations in organic synthesis. While alcohol dehydrogenases often perform this task very efficiently [3,4], mild homogeneous catalysts are scarce. Traditionally, these reactions have been performed with stoichiometric chromium reagents. However, today's environmental restrictions render most of the stoichiometric metal oxidants obsolete. For this purpose, the use of *tert*-butylhydroperoxide (TBHP) as an oxidizing agent has been investigated in conjunction with several homogeneous transition-metal catalysts, including chromium, ruthenium, rhodium, and copper [9–13].

The present Communication outlines our efforts to create active artificial metalloenzymes for the oxidation of alcohols using biotinylated transition metal complexes in the presence of (strept)avidin as host protein.

In contrast to transition-metal catalyzed hydrogenation reactions (where nearly any rhodium–diphosphine moiety is an active catalyst), the number of versatile

* Corresponding author. Tel.: +41 32 718 2516; fax: +41 32 718 2511.
E-mail address: thomas.ward@unine.ch (T.R. Ward).

Table 1
Oxidation of alcohols catalyzed by different artificial metalloenzymes^a

Entry	Coenzyme	Substrate	Host protein	Conversion (%) ^b
1	1	Phenethyl alcohol	Streptavidin	81
2	1	Benzyl alcohol	Streptavidin	68
3 ^c	1	Cyclohexanol	Streptavidin	80
4 ^d	1	Phenethyl alcohol	Streptavidin	43
5	2	Phenethyl alcohol	Streptavidin	69
6 ^c	3	Phenethyl alcohol	Streptavidin	3
7 ^c	4	Phenethyl alcohol	Streptavidin	12
8	1	Phenethyl alcohol	Ser112Gly	75
9	1	Phenethyl alcohol	Pro64Gly	70
10	1	Phenethyl alcohol	Avidin	92

^a Unless otherwise stated, the reaction was carried out under nitrogen at room temperature for 90 h with alcohol (62.5 μmol), aqueous TBHP (75.0 μmol), coenzyme (0.25 μmol) dissolved in DMF (31 μl), protein (0.08 μmol) dissolved in a mixture of water (500 μl) and acetone (100 μl).

^b Determined by GC integration.

^c The reaction was performed for 140 h.

^d The coenzyme was dissolved in DMSO (31 μl).




Fig. 1. Non-denaturing gel-electrophoresis demonstrating the integrity of the host protein following a catalytic run. (Lane 1: denatured pure streptavidin, revealing monomeric protein; Lane 2: non-denatured pure streptavidin, revealing tetrameric and oligotetrameric protein; Lane 3: aqueous reaction mixture resulting from a catalytic run, identical to Lane 2; Lanes 4 and 5: supernatant and precipitate, respectively, resulting from centrifugation at 16000g of a catalytic run).

benzoic acid in a ratio of 4:1 (68% total conversion (Table 1, Entry 2)).

With cyclic alcohols such as cyclohexanol (Table 1, Entry 3), the corresponding ketone was obtained without the formation of ring-opened or lactone products (80% conversion).

Using DMSO instead of DMF to solubilize the catalyst precursor, led to a decrease in conversion (Table 1, Entry 4). Dimethyl sulfoxide can act as a reductant in certain aerobic oxidation reactions catalyzed by late transition metals and dimethyl sulfone is produced during the reaction [18]. In contrast, DMSO is also a stoichiometric oxidant in a variety of chemical [19,20] and biological [21] oxidation reactions, yielding dimethyl sulfide as a byproduct. However, in our case there is no evidence for presence of either dimethyl sulfone or dimethyl sulphide (GC analysis). Presumably DMSO lowers the oxidation rate by coordinating to the Ru(II) center.

To optimize the catalytic activity of such metalloenzymes, two complementary approaches can be pursued: chemical modification of the first coordination sphere of the biotinylated catalyst or genetic modification of the protein.

Our first attempts focused on the use of a biotinylated bipyridine Ru(II) catalyst **2** (Table 1, Entry 5). With this coenzyme, a slight decrease in catalytic activity was observed (69% conversion, compared to 80% conversion for the sulfonamide Ru(II) complex **1**).

We also examined the oxidation of *sec*-phenethyl alcohol in the presence of $[\text{Rh}(\eta^5\text{-C}_5\text{Me}_5)(\text{Biot-Ligand})]$ **3** and $[\text{Ir}(\eta^5\text{-C}_5\text{Me}_5)(\text{Biot-Ligand})]$ **4**. The results displayed in Table 1 suggest that the iridium catalyst **4** was more active than its rhodium analog **3**. However, both systems are much less active than the corresponding ruthenium-based catalysts **1** and **2**: acetophenone was obtained in 3% yield and 12% yield after 140 h with catalyst precursors **3** and **4**, respectively (Table 1, Entries 6–7).

Having identified the most promising organometallic fragment (Table 1, Entry 1), we subjected the host protein to site-directed mutagenesis. Two streptavidin mutants were tested with the most promising organometallic catalyst precursor **1**. Neither Ser112Gly nor Pro64Gly streptavidin mutants displayed enhanced activity (Table 1, Entries 8–9). Using avidin as the host protein, we found that, in the presence of **1**, the oxidation of *sec*-phenethyl alcohol proceeds nearly to completion (Table 1, Entry 10).

In summary, we have developed *active* artificial metalloenzymes for the oxidation of alcohols in water based on the non-covalent incorporation of biotinylated d^6 piano stool complexes in (strept)avidin. Having established that the protein does not suffer from oxidative damage despite the harsh reaction conditions, our next goal is to develop *selective* artificial metallo-alcohol dehydrogenases for the kinetic resolution of secondary alcohols.

Acknowledgements

This work was supported by the Swiss National Science Foundation and CERC3. We thank Belovo Egg Science and Technology for a generous gift of egg white avidin.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.jorganchem.2005.02.001](https://doi.org/10.1016/j.jorganchem.2005.02.001).

References

- [1] E.N. Jacobsen, A. Pfaltz, H. Yamamoto (Eds.), *Comprehensive Asymmetric Catalysis*, vols. I–III, Springer, Berlin, 1999.
- [2] H.-U. Blaser, C. Malan, P. Pugin, F. Spindler, H. Steiner, M. Studer, *Adv. Synth. Catal.* 345 (2003) 103.
- [3] K. Faber, *Biotransformations in Organic Chemistry*, fifth ed., Springer, Berlin, 2004.
- [4] K. Drauz, H. Waldmann (Eds.), *Enzyme Catalysis in Organic Synthesis: a Comprehensive Handbook*, vols. I–II, VCH, Weinheim, 1995.
- [5] (a) J. Collot, J. Gradinaru, M. Skander, N. Humbert, A. Zocchi, T.R. Ward, *J. Am. Chem. Soc.* 125 (2003) 9030;
(b) M.T. Reetz, *Proc. Natl. Acad. Sci. USA* 101 (2004) 5716;
(c) M.E. Wilson, G.M. Whitesides, *J. Am. Chem. Soc.* 100 (1978) 306.
- [6] F. van de Velde, I.W.C.E. Arends, R.A. Sheldon, *J. Inorg. Biochem.* 80 (2000) 81.
- [7] M. Ohashi, T. Koshiyama, T. Ueno, M. Yanase, H. Fujii, Y. Watanabe, *Angew. Chem. Int. Ed.* 42 (2003) 1005.
- [8] J.R. Carey, S.K. Ma, T.D. Pfister, D.K. Garner, H.K. Kim, J.A. Abramite, Z. Whang, Z. Guo, Y. Lu, *J. Am. Chem. Soc.* 126 (2004) 10812.
- [9] K.B. Sharpless, T.R. Verhoeven, *Aldrichim. Acta* 12 (1979) 63.
- [10] J. Muzart, A. Nait-Ajjou, *J. Mol. Catal.* 66 (1991) 155.
- [11] J. Muzart, *Tetrahedron Lett.* 27 (1986) 3139.
- [12] S. Uemura, S.R. Patil, *Tetrahedron Lett.* 23 (1982) 4353.
- [13] C.-M. Che, K.-W. Cheng, M.C.W. Chan, T.-C. Lau, C.-K. Mak, *J. Org. Chem.* 65 (2000) 7996.
- [14] G.J. ten Brink, I.W.C.E. Arends, R.A. Sheldon, *Science* 287 (2000) 1636.
- [15] A. Nait Ajjou, *Tetrahedron Lett.* 42 (2001) 13.
- [16] K. Sato, M. Aoki, J. Takagi, R. Noyori, *J. Am. Chem. Soc.* 119 (1997) 12386.
- [17] G. Süß-Fink, B. Therrien, L. Vieille-Petit, M. Tschan, V.B. Romakh, T.R. Ward, M. Dadrás, G. Laurenczy, *J. Organomet. Chem.* 689 (2004) 1362, and references therein.
- [18] (a) J. Trocha-Grimshaw, H.B. Henbest, *Chem. Commun.* (1968) 1035;
(b) H. Gampp, D. Haspra, W. Spieler, A.D. Zuberbühler, *Helv. Chim. Acta* 67 (1984) 1019.
- [19] T.V. Lee, in: B.M. Trost, I. Fleming (Eds.), *Comprehensive Organic Synthesis*, vol. 7, Pergamon, New York, 1991, p. 291.
- [20] J.P. Caradonna, P.R. Reddy, R.H. Holm, *J. Am. Chem. Soc.* 110 (1988) 2139, and references therein.
- [21] R. Hille, *Chem. Rev.* 96 (1996) 2757.