

## TESTATE AMOEBAE (PROTISTS) AS PALAEOENVIRONMENTAL INDICATORS IN PEATLANDS

Mariusz LAMENTOWICZ<sup>1</sup>, Edward A. D. MITCHELL<sup>2</sup>

**Abstract.** Testate amoebae (or testaceans, *Testacea*, *Arcellaceans*) are unicellular eukaryotic organisms living in freshwater or most terrestrial habitats such as soils, mosses, lakes, rivers, as well as brackish habitats such as estuaries. They are very abundant in *Sphagnum* mosses, where they live in the top part of mosses and the oxygenated part of the peat. The tests (shells) of *Testacea* are well preserved in peat and to a lesser extent in lake sediments.

Efforts should be concentrated on constructing reliable regional transfer functions (mathematical representation of relation of species to environmental variables — presence of particular taxa in fossil material is the function of past environmental parameters), as they exist for some world areas, and there is a complete lack of them for central Europe. Polish data are exceptionally important because Poland is under several contrasted climatic influences — from oceanic to continental. The comparison of our data on climatic tendencies with those from Western Europe and the rest of the world will show how similar or different the responses of Polish peatlands might be.

The paper has three aims: (a) to present the need for ecological studies on testate amoebae in Central Europe, (b) to show the potential of reconstruction of past environment on the basis of multiproxy studies that include testate amoebae as an integral part of the palaeoecology toolbox and (c) to put our research efforts on testate amoebae in Poland in a more global perspective.

**Key words:** *Sphagnum*-dominated peatlands, testate amoebae, protozoa, transfer function, pollen analysis, Holocene, palaeoecology, hydrology, climate, Tuchola Pinewoods.

**Abstrakt:** Ameby skorupkowe (*Testacea*) to jednokomórkowe organizmy, które żyją w środowiskach słodkowodnych, w większości siedlisk lądowych (glebie, mszakach, jeziorach, rzekach) oraz w siedliskach słonawych (np. w ujściowych partiach rzek). *Testacea* występują licznie w mchach torfowcach (*Sphagnum*), gdzie zasiedlają ich górną część oraz natleniony stropowy fragment torfu. Skorupki ameb dobrze zachowują się w torfie oraz w mniejszym stopniu w osadach jeziornych.

Główny wysiłek powinien być skoncentrowany na budowaniu regionalnych funkcji transferu (transfer function; matematyczny obraz relacji organizmu w stosunku do środowiska - obecność poszczególnych taksonów w materiale kopalnym jest funkcją przeszłych parametrów środowiskowych). Dla kilku obszarów już one powstały ale brak tego rodzaju danych dla centralnej części Europy. Polskie dane są wyjątkowo istotne, ponieważ obszar ten znajduje się pod wpływem różnych wpływów klimatycznych — od klimatu oceanicznego do kontynentalnego. Porównanie naszych danych z informacjami z innych stanowisk w Europie i na świecie może wykazać w jak odmienny sposób reagują polskie torfowiska na zmiany klimatyczne.

Artykuł ma trzy cele: (a) zaprezentowanie potrzeby badań ekologii ameb skorupkowych w Europie centralnej, (b) przedstawienie potencjału rekonstrukcji palaeośrodowiska na bazie wielu wskaźników, gdzie ameby skorupkowe stanowią integralną część metod badań paleoekologii i (c) zaprezentowanie naszych badań nad *Testacea* w Polsce w szerszej, globalnej perspektywie.

**Słowa kluczowe:** torfowiska mszarne (*Sphagnum*), ameby skorupkowe, Protozoa, funkcja transferu, analiza pyłkowa, holocen, paleoekologia, hydrologia, klimat, Bory Tucholskie.

<sup>1</sup> Department of Biogeography and Palaeoecology, Adam Mickiewicz University, Fredry 10, 61-701 Poznan, Poland; e-mail: mariuszl@amu.edu.pl

<sup>2</sup> Laboratoire des Systèmes Écologique — ECOS — École Polytechnique Fédérale de Lausanne (EPFL), and Institut Fédéral de Recherches WSL, Antenne Romande, Case postale 96, CH-1015 Lausanne, Switzerland; e-mail: edward.mitchell@epfl.ch

## INTRODUCTION

Testate amoebae (or testaceans, *Testacea*, *Arcellaceans*) are unicellular eukaryotic organisms living in freshwater or most terrestrial habitats such as soils, mosses, lakes, rivers, as well as brackish habitats such as estuaries. They are very abundant in *Sphagnum* mosses, where they live in the top part of mosses and the underlying oxygenated part of the peat. The tests (shells) of *Testacea* are well preserved in peat and to a lesser extent in lake sediments. The abundance and distribution of species in mires is correlated to factors such as moisture, water table depth and pH for which many species have a limited ecological tolerance (stenotopy). This stenotopy makes it possible to use testate amoebae for palaeoenvironmental reconstruction (Charman, 2001) and indeed, they are now becoming increasingly popular in multi-proxy palaeoenvironmental studies of peatlands, especially for inferring moisture changes over the Holocene. Such studies can be used for land management and nature conservation purposes. Indeed, knowledge on the developmental history of a site and its responses to past natural and anthropogenic changes allows to assess how the ecosystem may respond to present and future changes (Scott *et al.*, 2001; Davis, Wilkinson, 2004). However, so far, ecological and palaeoecological investigation on testate amoebae are not very common in central Europe and are very rare in Poland.

In Poland, modern studies of testate amoebae from *Sphagnum* peatlands commenced in 2002 in the Department of Biogeography and Palaeoecology in Adam Mickiewicz University in Poznan in cooperation with Department of Biological Sciences of the University of Alaska Anchorage (former affiliation of E. Mitchell). The research began with the intention to combine palaeoecological (testate amoebae, pollen and plant macrofossil) methods used in mire studies.

This publication presents some potential uses of testate amoebae analysis in palaeoecological reconstructions. We present results of a palaeoecological study of two peatlands from NE Poland (Tuchola Pinewoods) (Lamentowicz, 2005) as well as results of an ecological investigation used to build two transfer functions (Lamentowicz, Mitchell, 2004). Currently, our efforts are concentrated on two particular factors — depth to the water table and pH, but further developments are possible. The paper has three aims: (a) to present the need for ecological studies on testate amoebae in Central Europe, (b) to show the potential of reconstruction of past environment on the basis of multiproxy studies that include testate amoebae as an integral part of the palaeoecology toolbox and (c) to put our research efforts on testate amoebae in Poland in a more global perspective.

## ECOLOGICAL STUDIES

Testate amoebae are generally regarded as soil organisms (Foissner, 1987). They are highly diverse in the acid, oligotrophic habitat represented by *Sphagnum* mosses growing on peatlands where they represent one of the main components of the soil microbial loop (Gilbert *et al.*, 1998).

In *Sphagnum* mires, living testate amoebae are restricted to top (living) part of the mosses (Fig. 1) and the underlying dead part. In the top part (1–5 cm long) the testate amoebae community is composed of both mixotrophic taxa (e.g. *Hyalosphenia papilo*, *Amphitrema flavum* or *A. wrightianum*) that contain endosymbiotic photosynthetic algae and heterotrophic taxa (Meisterfeld, 1977; Mitchell, Gilbert, 2004). Below that level, only heterotrophic species occurs and the assemblage is dominated by species such as *Nebela* sp. sp. and *Hyalosphenia elegans* (Schönborn, 1963; Grospietsch, 1990). The highest diversity is found in the transition zone between the upper and lower part of *Sphagnum* shoots (Meisterfeld, 1977). Under the living *Testacea* zone empty tests of dead individuals remain. Those necrocenoses are the main area of interest of palaeoecology (Fig. 1 A).

Knowledge on the distribution of species in microhabitats and the relationship between species assemblages and environmental factors is a prerequisite for future studies of soil ecology and palaeoecology. Each taxonomic group responds to more or less specific environmental gradients. Once the present ecological preferences of the species are known, the link between the present and the past is achieved using transfer functions (Birks, 1995).

For testate amoebae, the best-studied factors, and those for which the highest correlation are usually found, are moisture and water table depth. *Testacea* are very sensitive to moisture changes and this factor is also one of the most important in peatland ecology. Ombrotrophic bogs depend solely on precipitation for their water supply and the relative importance of precipitation versus evaporation has a profound influence on the structure and functioning of the ecosystem. Climate therefore influences the micro-topography, and also the composition of plant, animal and microbial communities. This link between climate and the structure and functioning of the system make ombrotrophic bogs especially sensitive to climate change (Charman, 2001, 2002 b; Charman *et al.*, 2001; Wilmshurst *et al.*, 2003; Charman *et al.*, 2004).

Another important factor is pH, which was we also modeled. But because pH is very variable not only during the year but also daily (Tahvanainen, Tuomaala, 2003), the interpretation is more difficult than for water table depth. Thus pH is less extensively discussed in the literature on *Testacea* and further studies are required to better understand the relationship between testate amoebae and this factor. The scarcity of extensive studies, covering the entire range of pH encountered in peatland is partly due to the natural rarity of transitional situations between highly acidic and circum-neutral conditions resulting in a bimodal distribution of pH in peatlands (Sjörs, Gunnarson, 2002; Tahvanainen, Tuomaala, 2003).

Testate amoebae are very sensitive to habitat changes and the heterogeneity of the environment, but are nevertheless



cal factors are assumed (Birks, 1995). The variable having the largest influence was water table depth (DWT) (sampled sites spanned -3 to 40 cm of DWT), which explained the majority of variation (21%). Taxa preferring wet habitat are e.g.: *Nebela carinata* and *Amphitrema wrightianum*, whereas on the opposite side of the RDA biplot are projected "dry" taxa e.g.: *Nebela militaris* and *Trigonopyxis arcua*. It should be also mentioned that *Assulina muscorum* is described in this study as a species of dry hummock. Our results are in agreement with previous studies (Warner, 1987; Tolonen *et al.*, 1992; Charman, 1997; Charman, Warner, 1997; Mitchell *et al.*, 2000 b; Booth, 2001, 2002), in which hydrology (usually as moisture or water table depth) was also the most important factor governing the distribution of testate amoebae. Currently, our model allows to reconstructing water table depth with an accuracy 9.89 cm (RMSEP) represented by weighted averaging model with tolerance downweighting of rare species. In the future, comparisons with other statistical models e.g. WA-PLS or PLS (Birks, 1995, 1998) are planned.

## PALAEOECOLOGICAL STUDIES

Testate amoebae have been used in ecological and palaeoecological studies since the beginning of the 20<sup>th</sup> century. Frey (1964) describes the development of early research on *Testacea*, in which Lagerheim (1901) was the first who recorded the tests in freshwater sediment. Steinecke (1914) was the first to recognize and describe the presence of rhizopods in *Sphagnum* habitat. Hesmer (1929) interpreted the lack of some testate amoebae at some depth in peat profiles as the result of test decomposition, and not of past changes in ecological conditions. Granlund (1932) related the degree of peat decomposition to testate amoebae taxa assemblages. Harnisch (1927, 1951) started working on peat cores, and consequently described testate amoebae from interglacial peat. Following semiquantitative methods of palaeoenvironmental reconstruction developed by Tolonen (1966), quantitative methods were applied (Woodland *et al.*, 1998; Wilmshurst *et al.*, 2002; Hendon, Charman, 2004) along with the development of multivariate statistics (Birks, 1995, 1998).

Testate amoebae (taxa such as *Assulina* sp. sp. and *Amphitrema flavum*) were sometimes counted along with pollen and their abundance related to the total pollen percentage (Aaby, Tauber, 1975; Aaby, 1976; Latałowa, 1999). Charman *et al.* (2000) pointed out that although this may be convenient it is not logical because "there is no ecological process that connects pollen influx with the living conditions of testate amoebae". The good representation of *Assulina muscorum* in pollen preparations is due to the high durability of its test (Charman, 2001). However this taxon, which is often regarded as a dry indicator by palynologists, in fact represents at least three distinct species — *Assulina muscorum*, *A. seminulum*, and *A. scandinavica*, each with its own ecological preferences (in this sequence, a decreasing tolerance to dryness with increasing size). These differences were revealed in a number of studies outside Poland (Warner, 1987; Tolonen *et al.*, 1992, 1994; Mitchell *et al.*, 2000 b; Booth, 2001) as well as in our Polish data (Lamentowicz, Mitchell, 2004). The low number of testate amoebae species in pollen

The range of acidity in the sites we sampled ranged between 3.26 and 6.67. As mentioned above, this factor is difficult to interpret and changes daily and annually. In the Tuchola Pine-woods area, some *Testacea* species occur in circum-neutral habitats e.g. *Quadrurella symmetrica* (idiosomic taxa) or *Centropyxis ecornis* (idiosomic pseudochitinous taxa) and some exclusively in acid habitats: *Nebela militaris* and *N. griseola* (both xenosomic taxa). The results for this factor also relate well to the other world studies, despite an imbalance in the pH range covered. Further investigations are required to fill the gap in transitional habitats. The last factor modeled was conductivity, but it was not statistically significant and was highly correlated with pH.

Our training set from Poland is expanding along with the predictive power of the transfer function. We gathered more than 60 samples from different habitats that will be used in future studies from Pomeranian *Sphagnum* dominated peatlands (Lamentowicz, Obremska, Mitchell in prep; Lamentowicz, Tobolski in prep). Further sampling in central Poland and mountain regions are also planned.

counts is explained by the pollen preparation method (acetolysis) of peat samples. This treatment destroys most tests, especially the agglutinated (e.g. *Diffflugia*) or siliceous (e.g. *Euglypha*) taxa (Hendon, Charman, 1997).

Multiproxy palaeoclimatic reconstruction studies in peatland are rare in Poland. Past changes in water level have mostly been obtained by using the peat humification index (Żurek, Pazdur, 2000; Żurek, Michczyńska *et al.*, 2002). In Poland there is presently a lack of comprehensive studies, comprising many bioindicators (Charman *et al.*, 1999; McCarroll *et al.*, 2003). The multiproxy approach allows reconstructing a much wider range of environmental changes in the Holocene (Barber *et al.*, 1998; Mauquoy, Barber, 2002; Mauquoy *et al.*, 2004). Testate amoebae were used similarly in some studies (Charman *et al.*, 1999, 2001; Mitchell *et al.*, 2001; Langdon *et al.*, 2003). Changes in peat accumulation rates and the communities of organisms living at the surface of peatlands are the key indicators in palaeoclimatic research.

Recent publications of Obremska and Lamentowicz (2002) and Lamentowicz (2005) represent pilot studies in Polish peatland research in which, in addition to standard palaeoecological methods (palynological and macrofossils analysis), an analysis of testate amoebae was applied. Polish material was analyzed qualitatively (Lamentowicz, 2005) and was compared with other studies. Two cores from two *Sphagnum* mires were studied. Figure 1 (B and C) presents the results of analysis of peat monolith from Jelenia Wyspa *Sphagnum* sampled in high resolution. Stratigraphic changes in sediment are noticeable and they are also reflected by testate amoebae taxa composition. The lower part of the monolith is characterized by the occurrence of *Amphitrema flavum* — species recorded in wet peat moss habitats. The testate amoebae data shows a gradual drying of the peatland surface connected with the onset of *Sphagnum* domination (Lamentowicz, 2003 & in prep.). The drying trend is also indicated by

the appearance of *Heleopera sylvatica*, *Nebela militaris* and *Nebela griseola* and the disappearance of *Amphitrema flavum*. This short core thus gives much information about past local hydrological conditions. Recently, Jelenia Wyspa bog has been dominated by *Sphagnum* species and its shift to ombrotrophy or oligotrophy should be regarded as local, therefore we probably cannot interpret amoebae changes in the context of regional palaeohydrology. Although non-ombrotrophic peatlands may also be useful for palaeoclimatic reconstructions (Booth *et al.*, 2004), strictly ombrotrophic bogs nevertheless give a better palaeoclimatological signal because they are dependent exclusively on rainfall (Charman, 2001). The analysed core from Tuchola kettle-hole bog (Tuchola Pine-woods, N Poland) reveals a connection between changes in

testate amoebae and vegetation inferred from pollen and macrofossils analyses (Lamentowicz, 2004). In this study three palaeoecological methods were used: testate amoebae, pollen and plant macrofossils analyses. The resulting data will be presented in a wider regional context (Lamentowicz, Obremska, Mitchell in prep). By combining several methods a more accurate picture of past environmental changes can be obtained and regional climate changes can be separated from the global signal. A critical aspect of this research is the use of dendrochronological methods (Krapiec, 1998; Cedro, 2003; Martinelli, 2004) that allow high precision temporal calibration of the signals obtained through other methods and can be correlated with peat stratigraphy (Gunnarson *et al.*, 2003; McCarroll *et al.*, 2003).

## SUMMARY AND CONCLUSIONS

Peatlands represent the most significant terrestrial carbon sink (Borren *et al.*, 2004; Grace, 2004), are very sensitive to climate changes, and have recorded past climatic fluctuations since the beginning of the Holocene (Blackford, 1993; Chambers, Charman, 2004).

Currently, there are only few multi-proxy studies of bogs in Poland. Most of the research is based on one or two methods, usually represented by the analysis of diatoms, cladocerans, plant macrofossils or pollen and spores. No attempt has yet been made to correlate several methods (e.g. testate amoebae, palynology, plant macrofossils and dendrochronology). Consequently, the testate amoebae method has been little used in Poland (Offierska, 1978, 1984; Offierska, Okuniewska, 1986; Offierska-Wawrzyniak, 1993).

Although testate amoebae are mostly believed to be cosmopolitan, the total number of regions in which the ecology of peatland testate amoebae has been studied is relatively low. It therefore remains possible that in some regions the composition of the testate amoebae fauna, or the ecological preference of some species, may be different from those of better-studied regions. A locally produced transfer function is therefore needed, and to fill the existing gap in central Europe an effort has begun to build the required data set of Polish, modern testate amoebae (Lamentowicz, Mitchell, 2004). This data set now needs to be extended to improve the accuracy of the re-

construction and to extend the habitat types covered by the model. In view of the ongoing and predicted future change in climate, we need more high-resolution palaeoclimatological studies in order to gain a better understanding of the response of natural ecosystems to past climatic changes. This is particularly true in areas where few similar studies have been done until today. palaeomoisture data inferred from testate amoebae can be correlated with other signals (Charman, 2001, 2002 a). We have the opportunity to reconstruct temperature and moisture changes during the Holocene because of the high sensitivity of tree species and soil microbes to climatic conditions and atmospheric chemistry (especially CO<sub>2</sub> concentrations). This period is exceptionally significant in palaeoclimatic studies (Follard, Karl, 2001).

Charman (2001) stated that the knowledge on peatland testate amoeba is in its infancy in Quaternary science. Efforts should be concentrated on constructing reliable regional transfer functions, as they exist for some world areas, and there is a complete lack of them for central Europe. Polish data are exceptionally important because Poland is under several contrasted climatic influences — from oceanic to continental. The comparison of our data on climatic tendencies with those from Western Europe and the rest of the world will show how similar or different the responses of Polish peatlands might be (Charman, Hendon, 2000; Follard, Karl, 2001).

## REFERENCES

- AABY B., 1976 — Cyclic climatic variations in climate over the past 5500 yr reflected in raised bogs. *Nature*, **263**: 281–284.
- AABY B., TAUBER H., 1975 — Rates of peat formation in relation to degree of humification of local environment, as shown by studies of raised bog in Denmark. *Boreas*, **4**: 1–17.
- BARBER K.E., DUMAYNE-PEATY L., HUGHES P.D.M., MAUQUOY D., SCAIFE R., 1998 — Replicability and variability of the recent macrofossil and proxy-climate record from raised bogs: field stratigraphy and macrofossil data from Bolton Fell Moss and Walton Moss, Cumbria, England. *J. Quatern. Sc.*, **13**, 6: 515–528.
- BEYENS L., CHARDEZ D., 1994 — On the habitat specificity of the testate amoebae assemblages from Devon Island (NWT, Canadian Arctic), with the description of a new species: *Diffflugia ovalisina*. *Arch. Protistenkd.*, **144**: 137–142.
- BIRKS H.J.B., 1995 — Quantitative palaeoenvironmental reconstructions. In: Statistical modelling of Quaternary science data (eds. D. Maddy and J.S. Brew): 161–254. Quaternary Research Association. Technical Guide No. 5. Cambridge.
- BIRKS H.J.B., 1998 — Numerical tools in palaeolimnology — progress, potentialities, and problems. *J. Paleolimn.*, **20**: 307–332.

- BIRKS H.J.B., 2001 — Maximum likelihood environmental calibration and the computer program WACALIB — a correction. *J. Paleolimn.*, **25**: 111–115.
- BLACKFORD J., 1993 — Peat bogs as sources of proxy climatic data: past approaches and future research. In: Climate change and human impact on the landscape (ed. D.J. Charman): 47–56. Chapman and Hall.
- BOOTH R.K., 2001 — Ecology of testate amoebae (Protozoa) in two lake Superior coastal wetlands: implications for paleoecology and environmental monitoring. *Wetlands*, **21**, 4: 564–576.
- BOOTH R.K., 2002 — Testate amoebae as paleoindicators of surface-moisture changes on Michigan peatlands: modern ecology and hydrological calibration. *J. Paleolimn.*, **28**, 3: 329–348.
- BOOTH R.K., JACKSON S.T., GRAY C.E.D., 2004 — Paleoecology and high-resolution paleohydrology of a kettle peatland in upper Michigan. *Quatern. Res.*, **61**: 1–13.
- BORREN W., BLEUTEN W., LAPSHINA E.D., 2004 — Holocene peat and carbon accumulation rates in the southern taiga of western Siberia. *Quatern. Res.*, **61**: 42–51.
- CEDRO A. 2003. Zmiany klimatyczne na Pomorzu Zachodnim w swietle analizy sekwencji przyrostów rocznych sosny zwyczajnej, daglezi zielonej i rodzimych gatunków debów. Oficyna IN PLUS. Szczecin.
- CHAMBERS F.M., CHARMAN D.J., 2004 — Holocene environmental change: contributions from the peatland archive. *The Holocene*, **14**, 1: 1–6.
- CHARMAN D.J., 1997 — Modelling hydrological relationships of testate amoebae (Protozoa: Rhizopoda) on New Zealand peatlands. *J. The Royal Society New Zealand*, **27**, 4: 465–484.
- CHARMAN D.J., 2001 — Biostratigraphic and palaeoenvironmental applications of testate amoebae. *Quatern. Sc. Rev.*, **20**: 1753–1764.
- CHARMAN D.J., 2002 a — Climate reconstruction from peatlands. In: Combining climate proxies (ed. A.E.K. Ojala) First ESF-HOLIVAR workshop. European Science Foundation & Geological Survey of Finland. Helsinki.
- CHARMAN D.J., 2002 b — Peatlands and environmental change. Wiley & Sons. Chichester.
- CHARMAN D.J., WARNER B., 1997 — The ecology of testate amoebae (Protozoa: Rhizopoda) in oceanic peatlands in Newfoundland, Canada: Modelling hydrological relationships for paleoenvironmental reconstruction. *Ecosc.*, **4**, 4: 555–562.
- CHARMAN D.J., HENDON D., 2000 — Long-term changes in soil water tables over the past 4500 years: relationships with climate and north Atlantic atmospheric circulation and sea surface temperature. *Climatic Change*, **47**, 1–2: 45–59.
- CHARMAN D.J., HENDON D., PACKMAN S., 1999 — Multi-proxy surface wetness records from replicate cores on an ombrotrophic mire: implications for Holocene palaeoclimate records. *J. Quatern. Sc.*, **14**, 5: 451–463.
- CHARMAN D.J., HENDON D., WOODLAND W.A., 2000 — The identification of testate amoebae (Protozoa: Rhizopoda) in peats. Quaternary Research Association Technical Guide No. 9. London.
- CHARMAN D.J., CASELDINE C., BAKER A., GEAREY B., HATTON J., PROCTOR C., 2001 — Paleohydrological records from peat profiles speleothems in Sutherland, northwest Scotland. *Quatern. Res.*, **55**: 223–234.
- CHARMAN D.J., BROWN A.D., HENDON D., KAROFELD E., 2004 — Testing the relationship between Holocene peatland palaeoclimate reconstructions and instrumental data at two European sites. *Quatern. Sc. Rev.*, **23**: 137–143.
- DAVIS S.R., WILKINSON D.M., 2004 — The conservation management value of testate amoebae as "restoration" indicators: speculations based on two damaged raised mires in northwest England. *The Holocene*, **14**, 1: 135–143.
- FOISSNER W., 1987 — Soil protozoa: fundamental problems, ecological significance, adaptations in ciliates and testaceans, bioindicators, and guide to literature. *Progr. in Protistol.*, **2**: 69–212.
- FOISSNER W., 1996 — Soil protozoan diversity: the state of the art. *Mitt. Deut. Bodenk. Gesell.*, **81**: 219–220.
- FOISSNER W., 1999 — Soil protozoa as bioindicators: pros and cons, methods, diversity, representative examples. *Agric. Ecosyst. Environm.*, **74**: 95–112.
- FOLLARD C.K., KARL T.R., 2001 — Observed climate variability and change. In: Climate change 2001: The Scientific Basis (eds. J.T. Houghton *et al.*): 101–165. Cambridge University Press. Cambridge.
- FREY D.G. 1964. Remains of animals in Quaternary lake and bog sediments and their interpretation. *Arch. Hydrobiol.*, **2**: 114 pp.
- GILBERT D., AMBLARD C., BOURDIER G., FRANCEZ A.J., 1998 — The microbial loop at the surface of a peatland: structure, function, and impact of nutrient input. *Microb. Ecol.* **35**: 83–93.
- GRACE J., 2004 — Understanding and managing the global carbon cycle. *J. Ecol.*, **92**: 189–202.
- GRANLUND E., 1932 — De Svenska Högmossarnas Geologi, deras Bildningsbetingelser, Utvecklingshistoria och Utbredning jämte sambandet mellan Högmossbildning och Försumpning. *Arsb. Sverig. geol. Unders.*, Ser. C., **373**, 26: 1–193.
- GROSPIETSCH T., 1990 — Rhizopoden in der Moorforchung. Moor und Torfkunde. G. K. Stuttgart.
- GUNNARSON B.E., BORGMARK A., WASTEGÅRD S., 2003 — Holocene humidity fluctuations in Sweden inferred from dendrochronology and peat stratigraphy. *Boreas*, **32**: 347–360.
- HARNISCH O., 1927 — Einige Daten zur rezenten und fossilen testaceen Rhizopodenfauna der Sphagnen. *Arch. Hydrobiol.* **18**: 345–360.
- HARNISCH O., 1951 — Daten zur Gestaltung der ökologischen Valenz der sphagnicolen Rhizopoden im Verlauf des Postglazials. *Dtsch. zool. Z.*, **1**: 222–133.
- HENDON D., CHARMAN D.J., 1997 — The preparation of testate amoebae (Protozoa: Rhizopoda) samples from peat. *The Holocene*, **7**: 199–205.
- HENDON D., CHARMAN D.J., 2004 — High resolution peatland water table changes for the past 200 years: the influence of climate and implications for management. *The Holocene*, **14**: 125–134.
- HESMER H., 1929 — Mikrofossilien in Torfen. *Paläont. Z.*, **11**: 245–257.
- KRAPIEC M., 1998 — Oak dendrochronology of the Neoholocene in Poland. *Folia Quatern.*, **69**: 5–133.
- LAGERHEIM G., 1901 — Om lämningar af Rhizopoder, Heliozoer och Tintinnider i Svergies och Finlands lakustrina Kvartäraflringar. *Geol. Fören. Stockh. Förh.*, **23**: 469–520.
- LAMENTOWICZ M., 2003 - Zmiany roślinności i zespołów korzenionózek skorupkowych Testacea (Protozoa) w późnym Holocenie na torfowisku mszarnym Jelenia Wyspa w rezerwacie Bagna nad Stążką (Tucholski Park Krajobrazowy). Litologia, Stratygrafia i Geneza Utworów Czwartorzędowych: **5**: 231-242. Wydawnictwo Naukowe UAM. Poznań.
- LAMENTOWICZ M., 2005 - Origin and development of natural and seminatural peatlands in Tuchola Forest Inspectorate [in Polish with English Sum.]. Department of Biogeography and Palaeoecology. Bogucki Wydawnictwo Naukowe. Poznań.
- LAMENTOWICZ M., MITCHELL E. A. D., 2004 - The ecology of testate amoebae (Protists) in Sphagnum in relation to peatland ecology. *Microb. Ecol.*, **50**, 1: 48–63.
- LANGDON P. G., BARBER K.E., HUGHES P.D.M., 2003 — A 7500-year peat-based paleoclimatic reconstruction and evidence for an 1100-year cyclicity in bog surface wetness from Temple Hill Moss, Pentland Hills, southeast Scotland. *Quatern. Sc. Rev.*, **22**: 259–274.

- LATAŁOWA M., 1999 — Zmiany hydrologiczne wywołane przez czynnik klimatyczny i działalność człowieka zapisane w torfowisku Kołczewo i osadach dennych Jeziora Raczego. *In: Ewolucja geosystemów nadmorskich południowego Bałtyku* (eds: R.K. Borówka *et al.*): 99–103. Bogucki Wyd. Nauk. Poznań-Szczecin.
- LINE J.M., BIRKS H.J.B., 1990 — WACALIB version 2.1 — a computer program to reconstruct environmental variables from fossil assemblages by weighted averaging. *J. Paleolimn.*, **3**, 3: 170–174.
- LINE J.M., TER BRAAK C.J.F., BIRKS H.J.B., 1994 — WACALIB version 3.3 — a computer program to reconstruct environmental variables from fossil assemblages by weighted averaging and to derive sample specific errors of prediction. *J. Paleolimn.*, **10**: 147–152.
- MARTINELLI N., 2004 — Climate from dendrochronology: latest developments and results. *Glob. Planet. Change*, **40**: 129–139.
- MAUQUOY D., BARBER K.E., 2002 — Testing the sensitivity of the palaeoclimatic signal from ombrotrophic peat bogs in the northern England and the Scottish Borders. *Rev. Palaeobot. Palynol.*, **119**: 219–240.
- MAUQUOY D., BLAAUW M., GEEL van B., BORROMEI A., QUATTROCCHIO M., CHAMBERS F.M., POSSNERT G., 2004 — Late Holocene climatic changes in Tierra del Fuego based on multiproxy analyses of peat deposits. *Quatern. Res.*, **61**: 148–158.
- MCCARROLL D., JALKANEN R., HICKS S., TUOVINEN M., GAGEN M., PAWELLEK F., ECKSTEIN D., SCHMITT U., AUTIO J., HEIKKINEN O., 2003 — Multiproxy dendroclimatology: a pilot study in northern Finland. *The Holocene*, **13**, 6: 829–838.
- MEISTERFELD R., 1977 — Die horizontale und vertikale Verteilung der Testaceen (Rhizopoda, Testacea) in Sphagnum. *Arch. Hydrobiol.*, **79**, 3: 319–356.
- MEISTERFELD R., 1997 — Thekoamöben — ihr Potential für Ökosystemforschung und Bioindikation. *Abh. Ber. Naturkundemus.*, **69**, 2: 87–95.
- MITCHELL E. A.D., GILBERT D., 2004 — Vertical micro-distribution and response to nitrogen deposition of testate amoebae in *Sphagnum*. *J. Eukar. Microbiol.*, **51**, 4: 485–495.
- MITCHELL E. A.D., BUTTLER A., WARNER B.G., GOBAT J.M., 1999 — Ecology of testate amoebae (Protozoa: Rhizopoda) in *Sphagnum*-dominated peatlands in the Jura mountains, Switzerland and France. *Ecosc.*, **6**, 4: 565–576.
- MITCHELL E.A.D., BORCARD D., BUTTLER A.J., GROSVERNIER P., GILBERT D., GOBAT J.M., 2000 a — Horizontal distribution patterns of testate amoebae (Protozoa) in *Sphagnum magellanicum* carpet. *Microb. Ecol.*, **39**: 290–300.
- MITCHELL E.A.D., BUTTLER A., GROSVERNIER P., RYDIN H., ALBINSSON C., GEENUP A. L., HEIJMANS M.M.P.D., HOOSBEEK M.R., SAARINEN T., 2000 b — Relationships among testate amoebae (Protozoa), vegetation and water chemistry in five *Sphagnum*-dominated peatlands in Europe. *New Phytologist*, **145**: 95–106.
- MITCHELL E.A.D., VANDER KNAPP W.O., LEEUWEN J.F.N., BUTTLER A., WARNER B.G., GOBAT J.M., 2001 — The palaeoecological history of the Praz-Rodet bog (Swiss Jura) based on pollen, plant macrofossils and testate amoebae (Protozoa). *The Holocene*, **11**, 1: 65–80.
- NGUYEN H., GILBERT D., BERNARD N., MITCHELL E. A.D., BADOT P.M., 2004 — Relationship between atmospheric pollution characterized by NO<sub>2</sub> concentrations and testate amoebae abundance and diversity. *Acta Protozool.*, **43**, 3: 233–240.
- OBREMSKA M., LAMENTOWICZ M., 2002 — Geology and history of kettle-hole bog on Brda outwash plain on the basis of pollen and testate amoebae (Protozoa) analyses. *In: Tuchola National Park in the concept of Biosphere Reserve* (eds: J. Banaszak and K. Tobolski) [in Polish with English Sum.]: 205–221. Charzykowy. Tuchola National Park.
- OFFIERSKA J., 1978 — Pelzaki skorupkowe (Testacea) torfowiska otaczającego j. Skrzynke w Wielkopolskim Parku Narodowym. *Bad. Fizj. Pol. Zach. Ser. C*, **31**: 7–38.
- OFFIERSKA J., 1984 — Zmienność skorupki u niektórych gatunków pelzaków skorupkowych (Testacea) w Wielkopolskim Parku Narodowym. *Bad. Fizj. Pol. Zach. Ser. C*, **34**: 154–162.
- OFFIERSKA J., OKUNIEWSKA I., 1986 — Analiza palinologiczna i fauny Testacea (Rhizopoda) z torfowiska przy jeziorze Skrzynka w Wielkopolskim Parku Narodowym. *Spraw. Pozn. TPN*, **103**: 146–148.
- OFFIERSKA-WAWRZYŃIAK J., 1993 — Kopalne i współczesne korzenionózki skorupkowe (Testacea: Rhizopoda) torfowiska Skrzynka w Wielkopolskim Parku Narodowym. *Bad. Fizj. Pol. Zach. Ser. C*, **31**: 5–20.
- SCHÖNBORN W., 1963 — Die Stratigraphie lebender Testaceen im Sphagnetum der Hochmoore. *Limnologica*, **1**, 4: 315–321.
- SCOTT D.B., MEDIOLI F.S., SCHAFFER C.T., 2001 — Monitoring in Coastal Environments Using Foraminifera and Thecoamoebian Indicators. Cambridge University Press. Cambridge.
- SJÖRS H., GUNNARSSON U., 2002 — Calcium and pH in north and central Swedish mire waters. *J. Ecol.*, **90**: 650–657.
- STEINECKE F., 1914 — Die beschalten Wurzelfüssler (Rhizopoda testacea) des Zehlaubruches. *Schr. phys. ökon. Ges. Königsb.*, **54**: 299–328.
- TAHVANAINEN T., TUOMAALA T., 2003 — The reliability of mire water pH measurements — a standard sampling: protocol and implications to ecological theory. *Wetlands*, **23**, 4: 701–708.
- TER BRAAK C.J.F., ŠMILAUER P., 1998 — CANOCO Reference Manual and User's Guide to Canoco for Windows Software for Canonical Community Ordination (version 4). Centre for Biometry. Wageningen.
- TOLONEN K., 1966 — Stratygraphic and rhizopod analyses of an old raised bog, Varrassuo, in Hollola, south Finland. *Ann. Botan. Fen.* **3**: 147–166.
- TOLONEN K., WARNER B.G., VASANDER H., 1992 — Ecology of Testaceans (Protozoa: Rhizopoda) in mires in southern Finland: I. Autecology. *Arch. Protistenk.*, **142**: 119–138.
- TOLONEN K., WARNER B.G., VASANDER H., 1994 — Ecology of Testaceans (Protozoa: Rhizopoda) in mires in southern Finland: II. Multivariate analysis. *Arch. Protistenk.*, **144**: 97–112.
- WARNER B.G., 1987 — Abundance and diversity of testate amoebae (Rhizopoda: Testacea) in *Sphagnum* peatlands in southwestern Ontario, Canada. *Arch. Protistenk.* **133**: 173–189.
- WILMSHURST J.M., MCGLONE M., CHARMAN D.J., 2002 — Holocene vegetation and climate change in southern New Zealand: linkages between forest composition and quantitative surface moisture reconstructions from an ombrotrophic bog. *J. Quatern. Sc.*, **17**, 7: 653–666.
- WILMSHURST J.M., WISER S.K., CHARMAN D.J., 2003 — Reconstructing Holocene water tables in New Zealand using testate amoebae: differential preservation of tests and implications for the use of transfer functions. *The Holocene*, **13**, 1: 61–72.
- WOODLAND W.A., CHARMAN D.J., SIMS P.C., 1998 — Quantitative estimates of water tables and soil moisture in Holocene peatlands from testate amoebae. *The Holocene*, **8**, 3: 261–273.