

Role of Soil Macrofauna in Phosphorus Cycling

Lydie Chapuis-Lardy, Renée-Claire Le Bayon, Michel Brossard, Danilo López-Hernández and Eric Blanchart

8.1 Introduction

Soils play a major role in the providing of goods and services by ecosystems to humans. They are involved in the main biogeochemical cycles (H₂O, C, N, P). They support agriculture, forestry, and pasture systems, and participate in climate regulation and detoxification. The importance of soils for ecosystem goods and services is based on their biological functioning, i.e., the whole biological functions carried out by soil biota in interactions with physical and chemical components of soil. These functions ensure, e.g., soil organic matter dynamics, nutrient recycling, soil structure and water retention. They are carried out by organisms of different sizes (from bacteria to macrofauna) and functional roles (decomposers, microbial regulators, soil engineers, and predators) (Lavelle et al. 2006).

Invertebrates of the macrofauna are key species for soil functioning. They participate in litter decomposition, mix organic and mineral matter, create and maintain soil structure by digging burrows and modifying aggregation, regulate microbial diversity and activity, and protect plants against pests and diseases (Lavelle et al. 2006). Following Jones et al. (1994), ecosystem engineers are

L. Chapuis-Lardy (✉)

Institut de Recherche pour le Développement (IRD), UMR 210 Eco&Sols, BP 1386, CP 18524, Dakar, Sénégal
e-mail: lydie.lardy@ird.fr

R.-C. Le Bayon

Soil and Vegetation Laboratory, Institute of Biology, University of Neuchâtel, Rue Emile-Argand 11, CP 158, 2009 Neuchâtel, Switzerland

M. Brossard and E. Blanchart

Institut de Recherche pour le Développement (IRD), UMR 210 Eco&Sols, Campus SupAgro, 2 place Viala, 34 060 Montpellier CEDEX 2, France

D. López-Hernández

Laboratorio de Estudios Ambientales, Instituto de Zoología y Ecología Tropical, Facultad de Ciencias, Universidad Central de Venezuela, Caracas, Venezuela

organisms able to modify the physical environment and availability of nutrients for other organisms. Lavelle (1997) defined the organisms fulfilling these functions in soil as soil engineers. They develop very complex interactions with other soil biota. For instance, the gut and biogenic structures (burrows, casts, nests) of earthworms and termites are specific habitats in which soil microbial activities are either stimulated or attenuated. Freshly egested earthworm casts are generally characterized by an intense mineralization of organic matter and the release of nutrients for plants. Conversely, the mineralization in old casts is reduced, which allows for long-term carbon and nutrient storage in soil (Martin 1991).

It is now recognized that soil engineers can improve soil fertility due to their role in nutrient dynamics at different spatial and temporal scales (review in Brown et al. 1999). They have been shown to affect the availability of nitrogen (N) and phosphorus (P), the main growth-limiting nutrients (e.g., Lee and Wood 1971; Bignell and Eggleton 2000; Brown et al. 2000; Brossard et al. 2007; Le Bayon and Milleret 2009). Numerous studies showed an increase in plant growth in the presence of earthworms, although the mechanisms are not fully understood; one of them being an increase in nutrient availability (Brown et al. 1999).

The objective of this chapter is to highlight, in a non-exhaustive view of published data, the role of soil macrofauna on P cycling. We will focus on the main soil engineers, i.e., earthworms and termites, whose roles in soil functioning have been frequently studied.

8.2 Earthworms

8.2.1 *Phosphorus Contents and Forms in Earthworm Biogenic Structures*

The first scientific studies on earthworm ecology, in the 1940s, related earthworm activity and soil fertility. Earthworms may affect soil structure and processes in different ways, according to their feeding and burrowing behaviors (synthesis in Brown et al. 2000). Three broad functional groups of earthworms have been described by Bouché (1977): epigeic, endogeic, and anecic. Epigeic earthworms live in the litter layer, consume plant litter and rarely ingest soil. Endogeics are mostly soil organic matter feeders and burrow extensively both horizontally and vertically within the soil. Anecics feed on particulate organic matter mixed with soil particles and often form deep, primarily vertical, burrows in which they bury surface litter. Numerous studies report results on chemical or physicochemical properties of earthworm casts, i.e., the by-products of gut passage, in comparison with surrounding soil. Most of these studies show that casts are characterized by higher P content, especially water-soluble or available (Bray- or Truog-P) P content (e.g., Nijhawan and Kanwar 1952; Bates 1960; Gupta and Sakal 1967; Graff 1970; Sharpley and Syers 1976; review in Edwards 1981; de Vleeschauwer and Lal 1981,

review in Lal 1987; Mulongoy and Bedoret 1989; López-Hernández et al. 1993). More recently, Kuczak et al. (2006) studied different P fractions (available P, moderately available P, and resistant P) in earthworm casts and soil in Amazonia. They observed that casts were characterized by higher total P and percentage of labile P pools (available and moderately available P). Guggenberger et al. (1996) found higher levels of alkali-extractable organic P in earthworm casts than in the surrounding oxisol. At this same site, Jiménez et al. (2003) worked in the field and analyzed inorganic and organic P fractions, including microbial P, in casts and surrounding soil. Under field conditions where earthworms ingest soil plus organic residues, total P concentrations were higher in casts than in surrounding soil. This increase was distributed over all inorganic and organic P fractions, including increased microbial P.

Because castings and earthworm-processed soils have the potential for higher P availabilities than uningested soils, the role of earthworm activities over the long term in soil P dynamics and the transfer of P into growing plants still needs consideration, especially in low-P status soils (e.g., Brown et al. 1999; Patrón et al. 1999; Chapuis-Lardy et al. 2009).

8.2.2 *Phosphorus Dynamics and Availability in Earthworm Casts*

Earthworm activity creates a series of geochemical and biological effects that are especially important for the cycling of P in soils (Brossard et al. 1996; Chapuis-Lardy et al. 1998, 2009; Nziguheba and Bünemann 2005; Le Bayon and Milleret 2009).

The availability of P is strongly influenced by the physical adsorption or fixation of P in soil (Frossard et al. 1995 and references therein). The phosphate ion (hereafter abbreviated as PO_4) concentration in the soil solution (C_P) may decrease as a direct consequence of P removal by root uptake. This depletion gives rise to a replenishment of P from the solid phase, which is time- and C_P -dependent. This PO_4 exchange reflects the buffer power of the soil for PO_4 and varies with the composition and the physicochemical conditions of the soil. Isotopic exchange kinetics (IEK) utilizing ^{32}P as a tracer has been extensively used to specifically assess inorganic P availability in a variety of soils (e.g., Morel et al. 1994; Fardeau 1996; Sinaj et al. 2001). Such approaches (see details in Frossard et al. 2011) have been used to evaluate P availability in earthworm casts or in soil in the presence or absence of earthworms.

We briefly focus here on the case of *Pontoscolex corethrurus* (Glossoscolecidae, Oligocheta), a widespread tropical endogeic earthworm. López-Hernández et al. (1993) reported higher P availability in fresh casts derived from two tropical soils with contrasting P sorption capacity. Chapuis and Brossard (1995) found higher C_P and $E_{(1 \text{ min})}$ (the amount of PO_4 that is isotopically exchanged within 1 min) values in casts egested by *P. corethrurus* fed with a Peruvian Ultisol. These changes in P availability were ascribed to (1) the higher pH of the gut content

(Barois and Lavelle 1986), (2) changes in sorption complexes induced by competition for sorbing sites between PO_4 and carboxyl groups of a mucus glycoprotein produced by the earthworm in its gut (López-Hernández et al. 1993), and (3) an increase in microbial activity during digestion (López-Hernández et al. 1993). More recently, IEK experiments were carried out to measure, describe, and compare the PO_4 exchange between liquid and solid phases of soil suspensions as a function of C_P and time, for two Malagasy soils and the associated casts egested by *P. corethrurus* (Chapuis-Lardy et al. 2009). The question arises whether the earthworm-induced modifications can change the ability of the soil solid phase to buffer C_P . The endogeic worm *P. corethrurus* increased the ability of PO_4 bound to the soil solid phase to exchange with PO_4 in soil solution (Fig. 8.1), because soil ingestion induced transformation of inorganic P to a more rapidly exchangeable form.

Tuffen et al. (2002) studied the transfer of ^{32}P in soil and between plants inoculated with arbuscular mycorrhizal fungi (AMF) in the presence of a temperate endogeic earthworm (*Aporrectodea caliginosa*). These authors reported enhanced ^{32}P transfer in soil and also between AMF-inoculated donor and receiver plants in the presence of earthworms.

Casts of earthworms usually contained larger amounts of organic P (e.g., for *P. corethrurus*, Chapuis-Lardy et al. 1998; Patrón et al. 1999), probably derived from a selective ingestion of soil particles (Chapuis-Lardy et al. 1998). Several




Fig. 8.1 Kinetics of isotopically exchangeable P ions (E , mg kg^{-1} soil) transferred between solid and liquid phases and the associated regression lines (E vs. time t) in control soil (from Andranomanelatra site) (crosses, dashed line) and casts of *Pontoscolex corethrurus* (triangles, semi-dashed line). The x - and y -axes are represented in log-log scales. Modified from Chapuis-Lardy et al. (2009)

studies reported increased enzymatic activity in earthworm casts and the stimulation of microbial activity (Sharples and Syers 1976; James 1991; López-Hernández et al. 1993; Chapuis and Brossard 1995; Brossard et al. 1996; Le Bayon and Binet 2006). However, Zhang et al. (2000) observed an increase of inorganic P in soils in the presence of earthworms, despite a decrease of acid and alkaline phosphatase activities in earthworm casts, suggesting that P-containing organic matter is digested in the gut rather than in the casts. This confirms the results of Devliegher and Verstraete (1996), who showed that the increase of inorganic P in soils in the presence of earthworms was due to the production of alkaline phosphatases by earthworms (Satchell and Martin 1984; Park et al. 1990) and a stimulation of microbial (acid) phosphatase production by earthworm activity (Satchell and Martin 1984). Devliegher and Verstraete (1996) concluded for the anecic earthworm *Lumbricus terrestris* that the increase of organic P mineralization and inorganic P availability in the presence of earthworms is more of a gut-associated process than a cast-associated process. Overall, the higher P availability observed in soils in the presence of earthworms may be linked to a rapid turnover of an increased organic P content. However, depending of the dominant species, earthworm activity could also result in the protection of organic P in stable macroaggregates and the increased fixation of inorganic P in Fe and Al hydroxides (Scheu and Parkinson 1994a, b; Suarez et al. 2004). It is concluded that earthworms markedly change the biogeochemical status of P (availability, organic P pool, enzymatic activities) in their guts and in the soil where they are active (drilosphere, sensu Lavelle et al. 1997), including casts and burrow linings. However, their impact on P dynamics and availability in the soil depends on the particular properties of soil, the organic P source, and the specific burrowing behavior and food preferences of worms.

The assumptions on earthworm-mediated transformations of organic P and related effects on P cycling in soils could be efficiently verified using soil labeling with radioactive P, especially if performed before chemical extraction (see methodological aspects in Frossard et al. 2011).

8.2.3 Surface-Cast Erosion and Phosphorus Transfer

The earthworm ecological category, and especially the feeding behavior, determines the type of surface-casts egested (Lee 1985; Curry and Schmidt 2007) as well as the architecture of the earthworm-induced macropores (Jégou et al. 2000; Bastardie et al. 2003). Such biogenic structures are involved in the regulation of soil physicochemical processes. However, how earthworms affect soil erosion is poorly understood (Blanchart et al. 2004). Surface-casts may be impacted by raindrops, and the fine soil particles they contain easily detached and transported during rainfall events (Le Bayon et al. 2002; Mariani et al. 2007). Structural stability of the biogenic structures is, together with rainfall intensity and field slope, an important determinant of soil erosion. Hence, earthworms can be proposed as important

moderators of soil erosion. Where stability is low and rainfall intense, material from the biogenic structures could be transported downstream by water runoff, thus leading to a significant loss of soil particles and associated P. Several studies have been performed to better understand the processes involved in surface-cast erosion and P transfer. Nevertheless, comparison between these investigations are difficult because of very variable experimental conditions (in situ annual monitoring or simulated rainfall events), different chemical analyses of P (colorimetric methods, anion exchange resins, sequential extractions, loss by ignition), and different expression of the results (actual measurements or estimations).

Sharpley and Syers (1976, 1977) and Sharpley et al. (1979) were the first to report a contribution of earthworm casts to P enrichment in runoff waters. They estimated under a temperate permanent pasture in New Zealand that surface-casts of the endogeic *A. caliginosa* may account for 45% of the annual transfer of P (0.25 kg of particulate P ha⁻¹, and 75% of the annual soil losses, i.e., 800 kg of soil ha⁻¹). In a tropical forest in Ivory Coast, the contribution of surface-casts to soil losses reached 1,200 kg ha⁻¹ year⁻¹, which corresponded to a maximum release of 0.23 kg of total P ha⁻¹ year⁻¹ (Nooren et al. 1995). Focusing on the impact of raindrops that cause the disintegration of fresh and dry old casts of *Martiodrilus carimaguensis* in Colombia, Mariani et al. (2007) used simulated rainfall events and demonstrated a significant effect of soil drying on cast dispersion. They estimated that the mass of fresh casts dispersed could range between 4 and 32 tons ha⁻¹ year⁻¹, which might be equivalent to 0.025 and 0.354 kg of available P ha⁻¹ year⁻¹ in savannah and pasture, respectively. However, opposite results were reported in a temperate agroecosystem in France by Le Bayon and Binet (2001). Rainstorm events were simulated on a maize plot with a gentle slope of 4.5% populated by two dominant species of earthworms, the endogeic *A. caliginosa* and the anecic *L. terrestris* (Binet and Le Bayon 1999). Despite a high cast abundance on the soil surface (25% of the area), amounts of P recovered in runoff waters were twice as high without surface-casts (0.35 and 0.86 kg of particulate P ha⁻¹ rainfall⁻¹ with and without casts, respectively). Earthworm casts were thus proposed to act as a physical brake on soil erosion by creating a surface roughness, and this was recently confirmed by Jouquet et al. (2008) in steep-slope ecosystems of North Vietnam. Nevertheless, Le Bayon and Binet (2001) observed that once the breaking-down point of the physical resistance of casts is reached, all surface-casts may then be quickly disintegrated and washed away. Hence, transfer of P can occur over a short distance through successive suspension/deposition of soil particles in the water runoff.

Simultaneous investigations were conducted on annual variations of earthworm surface-casting and P transfer (Le Bayon et al. 2002), including a 2-month period in spring when earthworm activities are enhanced (Le Bayon and Binet 1999). Working at such different temporal scales highlighted the possibility that the contribution of earthworm surface-casts to soil and P transfers could be over-estimated. For instance, a total of 0.05–0.20 kg of particulate P ha⁻¹ year⁻¹ were actually measured in runoff waters during one year, whereas 0.25–0.49 kg of particulate P ha⁻¹ year⁻¹ were estimated to be lost during the 2-month study of intensive earthworm activity in spring. Moreover, belowground egestions, which

may represent large amounts of casting activities (Decaëns et al. 1999; Bohlen et al. 2004), and earthworm burrows need to be taken into account for a better understanding of P fluxes (Jensen et al. 2000).

In conclusion, despite the high variability of the data presented in different studies, a contribution of earthworm casts and burrows to the transfer of P has been demonstrated. This is intimately linked both to biotic factors (earthworm species, ecological group, abundance, biomass, vegetation cover, etc.) and also abiotic parameters (slope, climate, intensity and frequency of the rainfall events, aggregate stability, etc.).

8.3 Termites

8.3.1 *Phosphorus Contents in Termite Mounds*

Termites play an important role in the transformation processes of organic compounds in savannahs and tropical forest ecosystems (e.g., Lee and Wood 1971; Bignell and Eggleton 2000; Holt and Lepage 2000; López-Hernández 2001). Data on the P content of termite mounds in relation to adjacent soils are sometimes contradictory. Extensive studies of mounds and other structures of various Australian and African termites showed little difference in P concentration between termite structures and the soil from which they were built (Lee and Wood 1971; Leprun and Roy-Noël 1977). However, Okello-Oloya et al. (1985) for Australian *Amitermes* mounds, and Nutting et al. (1987) for subterranean North American desert termites, found that extractable P was significantly greater in mounds than in associated soils. López-Hernández et al. (2006) reported, for neo-tropical termites, significantly higher levels of both total and available P in termite mounds of *Nasutitermes ephratae*, a common plant-debris-feeding termite, compared with the adjacent soil. Moreover, in a comprehensive P-fractionation study of *Nasutitermes*, López-Hernández (2001) showed that all P levels were significantly higher in the mounds than in the adjacent soils. Similar results were found when comparing $E_{(t)}$ (the amount of PO_4 that is isotopically exchanged within t minutes) values of isotopically exchangeable P (López-Hernández et al. 1989a). Rückamp et al. (2010) also reported P enrichment in various termite mounds collected from seven dominant Brazilian ecosystems.

The relative P enrichment of the mound may have two main sources: (1) a biological origin from salivary, fecal, and plant debris used in the mound building, and (2) a mineral origin from an accumulation of clay within the mound. As a result of different feeding habits, termites collect different materials to build their nest structures (López-Hernández et al. 2006). Wood et al. (1983) differentiate plant-debris-feeding termites, which employ the selection of certain particle sizes (e.g., clays with high P-fixing capacity from the subsoil) to cement soil particles with saliva alone or with saliva and feces (Lee and Wood 1971; Wood et al. 1983), from

soil-feeding termites that mainly employ ingestion of organic-rich fractions mixed with soil particles (mostly from the topsoil). Then, they reported that available P appears to be proportionally more abundant in mounds of soil-feeding termites than in the mounds of plant-debris-feeding termites when compared to adjacent soils.

8.3.2 Phosphorus Dynamics and Availability in Termite Mounds

Fardeau and Frossard (1992) showed that P isotopically exchangeable in 1 min [$E_{(1 \text{ min})}$] increased from 2.2 mg kg⁻¹ in the bulk soil to 87.0 mg kg⁻¹ in the centre of a *Trinervitermes geminatus* nest. The P sorption processes are strongly reduced in organic-matter-enriched mounds compared with adjacent soils for both grass-feeders *T. geminatus* and *N. ephratae* (Table 8.1). On the other hand, *Macrotermes* species forage mostly from the organic carbon-deficient subsoil and preferably selected finer soil particles for the nest construction (Leprun and Roy-Noël 1976; Pomeroy 1983; López-Hernández and Febres 1984; López-Hernández et al. 2006). Their mounds built with clay-enriched materials have very high P-sorbing capacities and consequently a lower P availability than the adjacent topsoil (Table 8.1). For the subterranean species (*Ancistrotermes cavithorax* and *Microtermes toumodiensis*), there was more P sorption in nest structures than in associated soils (Table 8.1). The low concentration of water-soluble P (C_P) found in the subsoil reflects strong P sorption in this soil layer, probably due to the presence of the materials enriched with high P-sorbing clay.

The higher levels of water-soluble inorganic P (C_P) in the termite nests presented in Table 8.1 may also result from the transformation of organic P through enzymatic activity in the fresh biostructures, as observed for earthworm casts (Sharpley and Syers 1976; Le Bayon and Binet 2006). Comparisons of phosphatase activities between termite nests and adjacent soils have already been made for several termite species (López-Hernández et al. 1989b; Roose-Amsaleg et al. 2005). Phosphatase activities in the soil and mounds for the South American *N. ephratae* ranged from 0.56 to 2.32 $\mu\text{mol } p\text{-nitrophenol g}^{-1} \text{ h}^{-1}$. Comparison of phosphatase activities for both mound and associated soils showed no statistical difference (López-Hernández

Table 8.1 Isotopically exchangeable P [$E_{(1 \text{ min})}$, mg kg⁻¹ soil] and water-soluble P (C_P , mg L⁻¹) in termite nests and adjacent soils

Termite species	Nest		Soil	
	$E_{(1 \text{ min})}$	C_P	$E_{(1 \text{ min})}$	C_P
<i>Macrotermes bellicosus</i>	0.29	0.002	0.34	0.021
<i>Trinervitermes geminatus</i>	3.11	0.221	0.19	0.010
<i>Cubitermes severus</i>	0.33	0.016	0.13	0.006
<i>Nasutitermes ephratae</i>	1.88	0.124	0.46	0.006
<i>Ancistrotermes cavithorax</i>	0.33	0.006	0.20	0.008
<i>Microtermes toumodiensis</i>	0.45	0.009	0.24	0.009

Adapted from López-Hernández et al. (2006)

et al. 1989a, b). Increases in enzyme activities and microbial biomass are expected to occur in mounds as a result of the increase in their organic matter content. Therefore, enzyme activity in the *N. ephratae* mounds might be inhibited as a consequence of the relatively high available P found in those structures. In contrast to the results of López-Hernández et al. (1989a), Roose-Amsaleg et al. (2005) found that soil-feeding termites affect soil phosphatase activity, with lower activities in mature nests than in surrounding soils. Comminution and mixing by termite activity allows for a greater surface area of substrate to come into contact with appropriate microorganisms and enzymes (Garnier-Sillam et al. 1987). Roose-Amsaleg et al. (2005) found that phosphatase activities tend to decrease with ageing from fresh to old and mature nests, and concluded that the released inorganic P could have a retroactive effect, inhibiting phosphatase activities as previously reported by López-Hernández et al. (1989a).

So far, it seems that termite feeding and nest-building habits have a strong influence on P availability parameters in mound, nest, and gallery structures. Humivorous and plant-feeding species cause a substantial increase of PO₄ bioavailability in nests, associated with a reduction of P sorption, whereas *Macrotermes* sp. (which in many cases build very large mounds) reduce PO₄ availability, with an associated increase of P sorption, by transporting finer soil materials from deeper soil layers.

8.3.3 *Termite Mounds and Phosphorus Transfer*

Phosphorus transport from the mound into the soil can happen in several ways. Termites might directly relocate nest materials into the soil. Furthermore, relocation can happen when termite mounds with dying colonies decay and erode. Although intact, inhabited termite mounds often have a dense surface that is particularly impermeable to water (e.g., Lal 1988; Contour-Ansel et al. 2000; Jouquet et al. 2004), the partial erosion and leaching from inhabited termite mounds may add P to the adjacent soil. However, erosion rates can be variable for a given species: Roose (1981) reported for *T. geminatus* that erosion of mounds can vary from 810 to 2,670 kg ha⁻¹ year⁻¹, representing 10–33% of the mound's mass (Brossard et al. 2007).

Coventry et al. (1988) measured the nutrient content of a range of termite mounds and used a lower estimated rate of erosion of the mounds to calculate the amounts of nutrients returned to soil each year. Given annual erosion losses from the mounds of 3.5% (Bonell et al. 1986), they estimated that mound-building termites are responsible for the return of at least 15 g ha⁻¹ of weak acid-extractable P per year. These estimates did not take into account the return of P as a result of erosion of structures other than mounds, such as feeding galleries. The amounts of soil contained in such structures may be of a similar magnitude to the amounts of soil incorporated in mounds (Lee and Wood 1971) and, if taken into account, might double the above estimates.

Finally, considering water infiltration as a major process, in particular in semi-arid and savannah areas, the influence of large macropores derived from soil macrofauna activity has been studied for about two decades (e.g., Elkins et al. 1986; Léonard et al. 2004). However, the results vary depending on measurement methods, and the processes involved deserve further consideration. Janeau and Valentin (1987) showed for *Trinervitermes* spp. that termite activity reduced the rate of water infiltration as a consequence of declines in soil surface porosity. However, Roose (1981), in a study on the sealing crusts in *T. geminatus*-affected soils, reported an increase in soil infiltration induced by termite activity. Léonard and Rajot (2001) showed that the influence of the large macropores made by termites was better described as a runoff interception process than as ponded infiltration. The P transfers through these processes (infiltration, runoff) are all related to P associated with soil particles. For soils with low-P status, the PO_4 pool in soil solution involved in these transfers is not well defined and very difficult to measure.

8.4 Conclusion

Macrofauna affects soil P cycling in a way that differs according to functional groups (i.e., feeding and construction behavior). However, P contents are usually larger in biogenic structures than in the surrounding soil, ultimately leading to higher P availability through a rapid turnover of organic P and/or enhanced PO_4 exchange. Further research efforts should be directed at investigating macrofauna-mediated processes in agricultural systems because these are potentially important for enhancing soil P availability to cultivated plants. Isotopic labeling techniques have the potential to elucidate soil P dynamics and should be extensively used to identify such processes. Controlled laboratory studies must be conducted to better understand the impact of gut passage, the microbial populations associated with guts, and the egested materials on P forms and availability in the biogenic structures. Experiments with a wide gamut of species and ecological strategies comprising a community will permit upscaling from the individual to community and population levels. Companion field studies should aim to better understand the volume of soil impacted and the dynamics of the biogenic structures produced, on a landscape scale. Coupled modeling approaches can also be useful to fully understand the complex interplay of biogeochemical interactions and to upscale from cast levels to soil profile and the entire ecosystem (Standing et al. 2007; Blanchart et al. 2009).

References

- Barois I, Lavelle P (1986) Changes in respiration rate and some physicochemical properties of a tropical soil during transit through *Pontoscolex corethrurus* (Glossoscolecidae, Oligochaeta). *Soil Biol Biochem* 18:539–541

- Bastardie F, Capowicz Y, de Dreuzy JR, Cluzeau D (2003) X-ray tomographic and hydraulic characterization of burrowing by three earthworm species in repacked soil cores. *Appl Soil Ecol* 24:3–16
- Bates JAR (1960) Studies on Nigerian forest soils. I. The distribution of organic matter in the profile and in various soil fractions. *J Soil Sci* 11:246–256
- Bignell DE, Eggleton P (2000) Termites in ecosystems. In: Abe T, Bignell DE, Higashi M (eds) *Termites: evolution, sociality, symbioses, ecology*. Kluwer, Dordrecht, pp 363–387
- Binet F, Le Bayon RC (1999) Space-time dynamics in situ of earthworm casts under temperate cultivated soils. *Soil Biol Biochem* 31:85–93
- Blanchart E, Albrecht A, Brown G, Decaëns T, Duboisset A, Lavelle P, Mariani L, Roose E (2004) Effects of tropical endogeic earthworms on soil erosion. *Agric Ecosyst Environ* 104:303–315
- Blanchart E, Marilleau N, Chotte JL, Drogoul A, Perrier E, Cambier C (2009) SWORM: an agent-based model to simulate the effect of earthworms on soil structure. *Eur J Soil Sci* 60:13–21
- Bohlen J, Parmelee RW, Blair JM (2004) Integrating the effects of earthworms on nutrient cycling across spatial and temporal scales. In: Edwards CA (ed) *Earthworm ecology*, 2nd edn. CRC, Boca Raton, pp 183–200
- Bonell M, Coventry RJ, Holt JA (1986) Erosion of termite mounds under natural rainfall in semi-arid tropical northeastern Australia. *Catena* 13:11–28
- Bouché MB (1977) Stratégies lombriciennes. *Ecol Bull* 25:122–132
- Brossard M, Lavelle P, Laurent JY (1996) Digestion of a vertisol by the endogeic earthworm *Polypheretima elongate*, Megascolecidae, increases soil phosphate extractability. *Eur J Soil Biol* 32:107–111
- Brossard M, López-Hernández D, Lepage M, Leprun JC (2007) Nutrient storage in soils and nests of mound-building *Trinervitermes* termites in Central Burkina Faso: consequences for soil fertility. *Biol Fertil Soils* 43:437–447
- Brown GG, Pashanasi B, Villenave C, Patron JC, Senapati BK, Giri S, Barois I, Lavelle P, Blanchart E, Blakemore RJ, Spain AV, Boyer J (1999) Effects of earthworms on plant production in the tropics. In: Lavelle P, Brussaard L, Hendrix P (eds) *Earthworm management in tropical agroecosystems*. CABI, Wallingford, pp 87–148
- Brown GG, Barois I, Lavelle P (2000) Regulation of soil organic matter dynamics and microbial activity in the drilosphere and the role of interactions with other edaphic functional domains. *Eur J Soil Biol* 26:177–198
- Chapuis L, Brossard M (1995) Modifications et stabilité du phosphore échangeable d'un ferralsol ingéré par un ver géophage. *CR Acad Sci II A* 320:587–592
- Chapuis-Lardy L, Brossard M, Lavelle P, Schouller E (1998) Phosphorus transformations in a Ferralsol through ingestion by *Pontoscolex corethrurus*, a geophagous earthworm. *Eur J Soil Biol* 34:61–67
- Chapuis-Lardy L, Ramianandroa RS, Randriamanantsoa L, Morel C, Rabeharisoa L, Blanchart E (2009) Modification of P availability by endogeic earthworms (Glossoscolecidae) in Ferralsols of the Malagasy Highlands. *Biol Fertil Soils* 45:415–422
- Contour-Ansel D, Garnier-Sillam E, Lachaux M, Croci V (2000) High performance liquid chromatography studies on the polysaccharides in the walls of the mounds of two species of termite in Senegal, *Cubitermes oculatus* and *Macrotermes subhyalinus*: their origin and contribution to structural stability. *Biol Fertil Soils* 31:508–516
- Coventry RJ, Holt JA, Sinclair DF (1988) Nutrient cycling by mound-building termites in low fertility soils of semi-arid tropical Australia. *Aust J Soil Res* 26:375–390
- Curry JP, Schmidt O (2007) The feeding ecology of earthworms – a review. *Pedobiologia* 50:463–477
- de Vleeschauwer D, Lal R (1981) Properties of worm casts in some tropical soils. *Soil Sci* 129:259–271
- Decaëns T, Rangel AF, Asakawa N, Thomas RJ (1999) Carbon and nitrogen dynamics in ageing earthworm casts in grasslands of the eastern plains of Colombia. *Biol Fertil Soils* 28:20–28

- Devliegher W, Verstraete W (1996) *Lumbricus terrestris* in a soil core experiment: effects of nutrient-enrichment processes (NEP) and gut-associated processes (GAP) on the availability of plant nutrients and heavy metals. *Soil Biol Biochem* 28:489–496
- Edwards CA (1981) Earthworms, soil fertility and plant growth. In: Apelhof M (ed) Proceedings of workshop on the role of earthworms in the stabilization of organic residues. Western Michigan University, Kalamazoo, pp 61–85
- Elkins N, Sabol G, Ward T, Whitford W (1986) The influence of subterranean termites on the hydrological characteristics of a Chihuahuan desert ecosystem. *Oecologia* 68:521–528
- Fardeau JC (1996) Dynamics of phosphate in soils – an isotopic outlook. *Fertil Res* 45:91–100
- Fardeau JC, Frossard E (1992) Processus de transformations du phosphore dans les sols d’Afrique de l’Ouest semi aride: application au phosphore assimilable. In: Tiessen H, Frossard E (eds) Phosphorus cycles in terrestrial and aquatic systems of semi arid Africa. SCOPE/UNEP proceedings. University of Saskatchewan, Saskatoon, pp 108–128
- Frossard E, Brossard M, Hedley MJ, Metherell A (1995) Reactions controlling the cycling of P in soils. In: Tiessen H (ed) Phosphorus in the global environment. Transfers, cycles and management – SCOPE 54. Wiley, New York, pp 107–137
- Frossard E, Achat DL, Bernasconi SM, Bünemann EK, Fardeau J-C, Jansa J, Morel C, Rabeharisoa L, Randriamanantsoa L, Sinaj S, Tamburini F, Oberson A (2011) The use of tracers to investigate phosphate cycling in soil–plant systems. In: Bünemann EK, Oberson A, Frossard E (eds) Phosphorus in action: biological processes in soil phosphorus cycling. *Soil biology*, vol 26. Springer, Heidelberg. doi:10.1007/978-3-642-15271-9_3
- Garnier-Sillam E, Villemeine G, Toutain F, Renoux J (1987) Contribution à l’étude du rôle des termites dans l’humification des sols forestiers tropicaux. In: Fedoroff N, Bresson JM, Courty MA (eds) Soil micromorphology. AFES, Plaisir, pp 331–335
- Graff O (1970) Phosphorus content in earthworm casts. *Landbauforsch Volk* 20:33–36
- Guggenberger G, Haumaier L, Thomas RJ, Zech W (1996) Assessing the organic phosphorous status of an Oxisol under tropical pastures following native savanna using ^{31}P NMR spectroscopy. *Biol Fertil Soils* 23:332–339
- Gupta ML, Sakal R (1967) The role of earthworms on the availability of nutrients in garden and cultivated soils. *J Indian Soc Soil Sci* 15:149–151
- Holt JA, Lepage M (2000) Termite and soil properties. In: Abe T, Bignell DE, Higashi M (eds) Termites: evolution, sociality, symbiosis, ecology. Kluwer, Dordrecht, pp 389–407
- James SW (1991) Soil, nitrogen, phosphorus, and organic matter processing by earthworms in tallgrass prairie. *Ecology* 72:2101–2109
- Janeau JL, Valentin C (1987) Relations entre les termitières *Trinervitermes* sp. et la surface du sol: réorganisations, ruissellement et érosion. *Rev Écol Biol Sol* 24:637–647
- Jégou D, Cluzeau D, Hallaire V, Balesdent J, Tréhen P (2000) Burrowing activity of the earthworms *Lumbricus terrestris* and *Aporrectodea giardi* and consequences on C transfers in soil. *Eur J Soil Biol* 36:27–34
- Jensen MB, Olsen TB, Hansen HCB, Magid J (2000) Dissolved and particulate phosphorus in leachate from structured soil amended with fresh cattle faeces. *Nutr Cycl Agroecosyst* 56:253–261
- Jiménez JJ, Cepeda A, Decaëns T, Oberson A, Friesen DK (2003) Phosphorus fractions and dynamics in surface earthworm casts under native and improved grasslands in a Colombian savanna oxisol. *Soil Biol Biochem* 35:715–727
- Jones CG, Lawton JH, Shachak M (1994) Organisms as ecosystem engineers. *Oikos* 69: 373–386
- Jouquet P, Tessier D, Lepage M (2004) The soil structural stability of termite nests: role of clays in *Macrotermes bellicosus* (Isoptera, Macrotermitinae) mound soils. *Eur J Soil Biol* 40:23–29
- Jouquet P, Podwojewski P, Bottinelli N, Mathieu J, Ricoy M, Orange D, Tran TT, Valentin C (2008) Above-ground earthworm casts affect water runoff and soil erosion in Northern Vietnam. *Catena* 74:13–21

- Kuczak CN, Fernandes ECM, Lehmann J, Rondon MA, Luizao FJ (2006) Inorganic and organic phosphorus pools in earthworm casts (Glossoscolecidae) and a Brazilian rainforest Oxisol. *Soil Biol Biochem* 38:553–560
- Lal R (1987) *Tropical ecology and physical edaphology*. Wiley, Chichester
- Lal R (1988) Effects of macrofauna on soil properties in tropical ecosystems. *Agric Ecosyst Environ* 24:101–116
- Lavelle P (1997) Faunal activities and soil processes: adaptive strategies that determine ecosystem function. *Adv Ecol Res* 27:93–132
- Lavelle P, Bignell DE, Lepage M, Volters V, Roger P, Ineson P, Heal W, Dillion S (1997) Soil function in a changing world: the role of invertebrate ecosystem engineers. *Eur J Soil Biol* 33:159–193
- Lavelle P, Decaëns T, Aubert M, Barot S, Blouin M, Bureau F, Margerie P, Mora P, Rossi JP (2006) Soil invertebrates and ecosystem services. *Eur J Soil Biol* 42:S3–S15
- Le Bayon RC, Binet F (1999) Rainfall effects on erosion of earthworm casts and phosphorus transfer by water runoff. *Biol Fertil Soils* 30:7–13
- Le Bayon RC, Binet F (2001) Earthworm surface casts affect soil erosion by runoff water and phosphorus transfer in a temperate maize crop. *Pedobiologia* 45:430–442
- Le Bayon RC, Binet F (2006) Earthworms change the distribution and availability of phosphorus in organic substrates. *Soil Biol Biochem* 38:235–246
- Le Bayon RC, Milleret R (2009) Effects of earthworms on phosphorus dynamics – a review. *Dyn Soil Dyn Plant* 3(2):21–27
- Le Bayon RC, Moreau S, Gascuel-Oudoux C, Binet F (2002) Annual variations in earthworm surface-casting activity and soil transport by water runoff under a temperate maize agroecosystem. *Geoderma* 106:121–135
- Lee KE (1985) *Earthworms, their ecology and relationships with soils and land use*. Academic, Bowen Hills
- Lee KE, Wood TG (1971) *Termites and soils*. Academic, London
- Léonard J, Rajot JL (2001) Influence of termites on runoff and infiltration: quantification and analysis. *Geoderma* 104:17–40
- Léonard J, Perrier E, Rajot JL (2004) Biological macropores effect on runoff and infiltration: a combined experimental and modelling approach. *Agric Ecosyst Environ* 104:277–285
- Leprun J-C, Roy-Noël J (1976) Minéralogie des argiles et répartition des nids épigés de deux espèces du genre *Macrotermes* au Sénégal occidental (Presqu'île du Cap-Vert). *Insectes Soc* 23:535–547
- Leprun J-C, Roy-Noël J (1977) Les caractères analytiques distinctifs des matériaux des nids du genre *Macrotermes* au Sénégal occidental. Leurs rapports avec les sols. *Pedobiologia* 17:361–368
- López-Hernández D (2001) Nutrient dynamics (C, N and P) in termite mounds of *Nasutitermes ephratae* from savannas of the Orinoco llanos (Venezuela). *Soil Biol Biochem* 33:747–753
- López-Hernández D, Febres A (1984) Changements chimiques et granulométriques produits dans des sols de Côte d'Ivoire par la présence de trois espèces de termites. *Rev Écol Biol Sol* 21:477–489
- López-Hernández D, Niño M, Fardeau JC, Nannipieri P, Chacón P (1989a) Phosphorus accumulation in savanna termite mound in Venezuela. *J Soil Sci* 40:635–640
- López-Hernández D, Niño M, Nannipieri P, Fardeau JC (1989b) Phosphatase activity in *Nasutitermes ephratae* termite nest. *Biol Fertil Soils* 7:134–137
- López-Hernández D, Lavelle P, Fardeau JC, Niño M (1993) Phosphorus transformations in two P-sorption contrasting tropical soils during transit through *Pontoscolex corethrus* (Glossoscolecidae, Oligochaeta). *Soil Biol Biochem* 25:789–792
- López-Hernández D, Brossard M, Fardeau JC, Lepage M (2006) Effect of different termite feeding groups on P sorption and P availability in African and South American savannas. *Biol Fertil Soils* 42:207–214

- Mariani L, Jiménez JJ, Torres EA, Amezcua E, Decaëns T (2007) Rainfall impact effects on ageing casts of a tropical anecic earthworm. *Eur J Soil Sci* 58:1525–1534
- Martin A (1991) Short- and long-term effects of the endogeic earthworm *Millsonia anomala* (Omodeo) (Megascolecidae, Oligochaeta) of tropical savannas on soil organic matter. *Biol Fertil Soils* 11:234–238
- Morel C, Tiessen H, Moir J, Stewart JWB (1994) Phosphorus transformations and availability due to crop rotations and mineral fertilization assessed by an isotopic exchange method. *Soil Sci Soc Am J* 58:1439–1445
- Mulongoy K, Bedoret A (1989) Properties of worm casts and surface soils under various plant covers in the humid tropics. *Soil Biol Biochem* 21:197–203
- Nijhawan SD, Kanwar JS (1952) Physicochemical properties of earthworm castings and their effect on the productivity of soil. *Indian J Agric Sci* 22:357–373
- Nooren CAM, van Breemen N, Stoorvogel JJ, Jongmans AG (1995) The role of earthworms in the formation of sandy surface soils in a tropical forest in Ivory Coast. *Geoderma* 65:135–148
- Nutting WL, Haverly MI, Lafage JP (1987) Physical and chemical alteration by two subterranean termite species in Sonoran Desert grassland. *J Arid Environ* 12:233–239
- Nziguheba G, Bünemann EK (2005) Organic phosphorus dynamics in tropical agroecosystems. In: Turner BL, Frossard E, Baldwin DS (eds) *Organic phosphorus in the environment*. CABI, Wallingford, pp 243–268
- Okello-Oloya T, Spain AV, John RD (1985) Selected chemical characteristics of the mounds of two species of *Amitermes* (Isoptera, Termitinae) and their adjacent surface soil from North-eastern Australia. *Rev Écol Biol Sol* 22:291–311
- Park SC, Smith TJ, Bisesi MS (1990) Hydrolysis of bis(4-nitro phenyl) phosphate by the earthworm *Lumbricus terrestris*. *Soil Biol Biochem* 22:729–730
- Patrón JC, Sánchez P, Brown GG, Brossard M, Barois I, Gutiérrez C (1999) Phosphorus in soil and *Brachiaria decumbens* plants as affected by the geophagous earthworm *Pontoscolex corethrurus* and P fertilization. *Pedobiologia* 43:547–556
- Pomeroy DE (1983) Some effects of mound building termites on soils of Uganda. *J Soil Sci* 27:377–394
- Roose E (1981) Dynamique actuelle de sols ferrallitiques et ferrugineux tropicaux d'Afrique occidentale. Étude expérimentale des transferts hydrologiques et biologiques de matières sous végétations naturelles ou cultivées. *Trav Doc ORSTOM* 130. ORSTOM, Paris
- Roose-Amsaleg C, Mora P, Harry M (2005) Physical, chemical and phosphatase activities characteristics in soil-feeding termite nests and tropical rainforest soils. *Soil Biol Biochem* 37:1910–1917
- Rückamp D, Amelung W, Theisz N, Bandeira AG, Martius C (2010) Phosphorus forms in Brazilian termite nests and soils: relevance of feeding guild and ecosystems. *Geoderma* 155:269–279
- Satchell JE, Martin K (1984) Phosphatase activity in earthworm faeces. *Soil Biol Biochem* 16:191–194
- Scheu S, Parkinson D (1994a) Effects of earthworms on nutrient dynamics, carbon turnover and microorganisms in soil from cool temperate forests of the Canadian Rocky Mountains: laboratory studies. *Appl Soil Ecol* 1:113–125
- Scheu S, Parkinson D (1994b) Effects of an invasion of an Aspen woodland (Alberta, Canada) by *Dendrobaena octaedra* on plant growth. *Ecology* 75:2348–2361
- Sharpley AN, Syers JK (1976) Potential role of earthworms casts for the phosphorus enrichment of run-off water. *Soil Biol Biochem* 8:341–346
- Sharpley AN, Syers JK (1977) Seasonal variation in casting activity and in the amounts and release to solution of phosphorous forms in earthworm casts. *Soil Biol Biochem* 9:227–231
- Sharpley AN, Syers JK, Springett JA (1979) Effect of surface-casting earthworms on the transport of phosphorus and nitrogen in surface runoff from pasture. *Soil Biol Biochem* 11:459–462

- Sinaj S, Buerkert A, El-Hajj G, Bationo A, Traore H, Frossard E (2001) Effects of fertility management strategies on phosphorus bioavailability in four West African soils. *Plant Soil* 233:71–83
- Standing D, Baggs EM, Wattenbach M, Smith P, Killham K (2007) Meeting the challenge of scaling up processes in the plant–soil–microbe system. *Biol Fertil Soils* 44:245–257
- Suarez ER, Fahey TJ, Groffman PM, Bohlen PJ, Fisk MC (2004) Effects of exotic earthworms on soil phosphorus cycling in two broadleaf temperate forests. *Ecosystems* 7:28–44
- Tuffen F, Eason WR, Scullion J (2002) The effect of earthworms and arbuscular mycorrhizal fungi on growth of and ^{32}P transfer between *Allium porrum* plants. *Soil Biol Biochem* 34:1027–1036
- Wood TG, Johnson RA, Anderson JM (1983) Modification of soils in Nigerian savanna by soil-feeding *Cubitermes*, Termitidae. *Soil Biol Biochem* 15:575–579
- Zhang BG, Li GT, Shen TS, Wang JK, Sun Z (2000) Changes in microbial biomass C, N and P and enzyme activities in soil incubated with the earthworms *Metaphire guillelmi* or *Eisenia fetida*. *Soil Biol Biochem* 32:2055–2062