

# Endosymbionts and honey bee colony losses?

Alexandre Aebi<sup>1</sup> and Peter Neumann<sup>2,3</sup>

<sup>1</sup> Agroscope Reckenholz-Tänikon, Research Station ART, Reckenholzstrasse 191, CH-8046 Zürich, Switzerland

<sup>2</sup> Swiss Bee Research Centre, Agroscope Liebefeld-Posieux, Research Station ALP, Schwarzenburgstrasse 161, CH-3003 Bern, Switzerland

<sup>3</sup> Department of Zoology and Entomology, Rhodes University, Grahamstown 6140, South Africa

Honey bees, *Apis mellifera*, are essential pollinators for the maintenance of natural biodiversity and agriculture [1]. Colony losses witnessed throughout the Northern hemisphere are therefore worrying [2], especially because no single driver has yet emerged as the definitive cause [3]. Interactions between viruses, ectoparasitic mites and microsporidian endoparasites are most likely key factors [3–5], but the underlying mechanisms are not well understood. Although it is known that maternally-inherited, facultative bacterial endosymbionts such as *Wolbachia* or *Rickettsia* can significantly interfere with viral and fungal infections of arthropods [6], they have so far been neglected in this regard. Here we propose to evaluate the potential role of such endosymbionts for colony losses.

Endosymbionts are widespread [7] in arthropods and transmitted vertically [8], but can only spread in host populations when infected females have a higher fitness, e.g. via providing protection against viruses or fungi [6]. For example, *Wolbachia* can protect the host against several vectored RNA viruses [9] and can be regarded as part of host immunity [6]. However, endosymbionts such as *Spiroplasma* and *Hamiltonella* can also be beneficial for their host's vectorial capacity, e.g. in the whitefly *Bemisia tabaci*–Tomato yellow leaf curl virus system, *Hamiltonella* protects viral particles in the vector [6,10].

To shed light on the potential influence of endosymbionts on losses, we here suggest an investigation of symbiont-mediated host protection against viruses transmitted by parasitic mites and/or associated with microsporidians (e.g. *Nosema ceranae*) [5], which could contribute to the tolerance of honey bee populations, e.g. against the mite *Varroa destructor* [11]. Moreover, endosymbionts carried by parasitic mites might favour virus transmission to and virulence in honeybees, which could explain regional differences in the impact of mites [11]. A combination of metagenomics and laboratory experiments appears suitable to compare the bacterial and viral communities associated with honey bees and their parasites in host populations

with or without elevated losses [2]. In conclusion, it seems as if endosymbionts play a role in honey bee pathology [12] and should therefore be investigated as a potential key to our understanding of major colony losses.

## Acknowledgements

We thank Vincent Dietemann, Einat Zchori-Fein, Daniele Daffonchio, Ameer Cherif and Renate Zindel for stimulating discussions and the FP7 projects STEP and BEEDOC (P.N.), the COST Actions FA0803 COLOSS (P.N. and A.A.) and FA0701 (A.A.) for financial support.

## References

- 1 Klein, A.M. *et al.* (2007) Importance of pollinators in changing landscapes for world crops. *Proc R. Soc. Lond. B.* 274, 303–313
- 2 Neumann, P. and Carreck, C. (2010) Honey bee colony losses: a global perspective. *J. Apic. Res.* 49, 1–6
- 3 Potts, S.G. *et al.* (2010) Global pollinator declines: drivers and impacts. *Trends Ecol. Evol.* 25, 345–353
- 4 Yang, X. and Cox-Foster, D.L. (2005) Impact of an ectoparasite on the immunity and pathology of an invertebrate: Evidence for host immunosuppression and viral amplification. *Proc. Natl. Acad. Sci. U.S.A.* 102, 7470–7475
- 5 Bailey, L. *et al.* (1983) Association of virus with two protozoal pathogens of the honey bee. *Ann. Appl. Biol.* 103, 13–20
- 6 Zindel, R. *et al.* (2011) Arthropod symbioses, a neglected parameter in pest- and disease-control programmes. *J. Appl. Ecol.* 48, 864–872
- 7 Hilgenboeker, K. *et al.* (2008) How many species are infected with *Wolbachia*? A statistical analysis on current data. *FEMS Microbiol. Lett.* 281, 215–220
- 8 O'Neill, S.L. *et al.* (1997) *Influential passengers: Inherited microorganisms and arthropod reproduction*, Oxford University Press
- 9 Teixeira, L. *et al.* (2008) The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol.* 6, e1000002
- 10 Gottlieb, Y. *et al.* (2010) The transmission efficiency of Tomato Yellow Leaf Curl Virus by the whitefly *Bemisia tabaci* is correlated with the presence of a specific symbiotic bacterium species. *J. Virol.* 84, 9310–9317
- 11 Rosenkranz, P. *et al.* (2010) The biology and control of *Varroa destructor*. *J. Inv. Pathol.* 103, 96–119
- 12 Evans, J.D. and Armstrong, T.N. (2006) Antagonistic interactions between honey bee bacterial symbionts and implications for disease. *BMC Ecol.* 6, DOI: 10.1186/1472-6785-6-4